

Commutation relations for SLE

Julien Dubédat*

1st February 2008

Abstract

Schramm-Loewner Evolutions (SLEs) describe a one-parameter family of growth processes in the plane that have particular conformal invariance properties. For instance, SLE can define simple random curves in a simply connected domain. In this paper we are interested in questions pertaining to the definition of several SLEs in a domain (i.e. several random curves). In particular, one derives infinitesimal commutation conditions, discuss some elementary solutions, study integrability conditions following from commutation and show how to lift these infinitesimal relations to global relations in simple cases. The situation in multiply-connected domains is also discussed.

For plane critical models of statistical physics, such as percolation or the Ising model, the general line of thinking of Conformal Field Theory leads to expect the existence of a non-degenerate scaling limit that satisfies conformal invariance properties. Though, it is not quite clear how to define this scaling limit and what conformal invariance exactly means.

One way to proceed is to consider a model in a, say, bounded (plane) simply- connected domain with Jordan boundary, and to set boundary conditions so as to force the existence of a macroscopic interface connecting two marked points on the boundary. In this set-up, Schramm has shown that the possible scaling limits satisfying conformal invariance along with a “domain Markov” property are classified by a single positive parameter $\kappa > 0$, in the seminal article [19]. This defines the family of Schramm-Loewner Evolutions (SLEs), that are probability measures supported on non-self-traversing curves connecting two marked boundary points in a simply-connected domain.

Consider the following situation for critical site percolation on the triangular lattice: a portion of the triangular lattice with mesh ε approximates a fixed simply connected domain D with two points x and y marked on the boundary. The boundary arc (xy) is set to blue and (yx) is set to yellow; sites are blue or yellow with probability $1/2$. Then the interface between blue sites connected to (xy) and yellow sites connected to (yx) is a non-self traversing curve from x to y . In this set-up, Smirnov has proved that the interface converges to SLE_6 , as conjectured earlier by Schramm ([21, 4]).

For discrete models such as percolation or the Ising model, the full information can be encoded as a collection of contours (interfaces between blue and yellow, $+$ and $-$ spins, ...). Hence it is quite natural to consider scaling limits as collection of contours, as in [1, 4]. Comparing the ideas of isolating one macroscopic interface by setting appropriate boundary conditions (following Schramm), and considering the scaling limit as a collection of contours, one is led to the problem of describing the joint law of a finite number of macroscopic interfaces created by appropriate boundary conditions.

For instance, for percolation, consider a simply connected domain with $2n$ marked points on the boundary, the $2n$ boundary arcs being alternatively blue and yellow. This gives n interfaces pairing the $2n$ points. One can then consider the joint scaling limits of these interfaces (either unconditionally or conditionally on a

*Courant Institute

given pairing). Each of these interfaces close to its starting point is absolutely continuous w.r.t. SLE_6 . So we are defining n “non-crossing” SLE_6 ’s; the problem is then to precisely quantify their interaction.

One remarkable feature of Schramm’s construction is the classification by a single positive parameter κ for one interface satisfying simple axioms. It is not hard to see that for, say, 2 interfaces connecting 4 points, each interface is a priori described by κ and a drift term materializing the interaction. This drift term can be seen as a function of the cross-ratio of the four boundary points.

The main goal of this article is to elucidate the constraints on the drift terms imposed by the general geometric framework, and to prove that in the most natural cases, the possible probability laws are characterized by a finite number of parameters. The geometric condition is that one can grow the interfaces in any order, at any relative speed, and get the same result in distribution.

We aim at defining several SLEs in the same simply-connected domain. As the growth of each SLE perturbs the time scales of other SLEs, we want the collection of SLE to be invariant in distribution under a global time reparametrization (that is, a time change $\mathbb{R}_+^n \rightarrow \mathbb{R}_+^n$). At an infinitesimal level, this invariance is expressed as a commutation relation for differential operators (the infinitesimal generators of the driving processes of the SLEs).

The conditions on the drift terms are non-linear differential equations involving the drift terms pairwise. If one writes the drift terms as log derivatives (in Girsanov fashion), then these conditions can be written as a system of linear PDEs of a certain form satisfied by a single “partition function”. The case where $2n$ points are marked on the boundary and n SLEs are grown is of particular interest. The system of PDEs satisfied by the partition function is then unique and the solution space is finite dimensional. The study of explicit solutions is the subject of the companion paper [10].

There is an analytically simple example of n SLEs in a domain with $(n + 1)$ marked points. Here the drift terms are rational (and SLEs are an example of $\text{SLE}_\kappa(\rho)$). In this case, we prove that infinitesimal commutation conditions can be lifted to global commutation; restriction formulae are derived for these. The radial analogue is also discussed.

For multiply connected domains, as for simply-connected domains with, say, more than 3 points marked on the boundary, the moduli space is no longer a point, so identifying the drift terms that give “physically relevant” SLEs is a problem. We work out the conditions imposed by the following intuitive criterion: the interface can be grown “from both ends”. This involves “cocycles” on the configuration space. The connection between this commutation condition and the restriction property framework ([13, 22]) is made explicit.

These questions are connected with work by Oded Schramm and David Wilson, Greg Lawler, and John Cardy; and also by Bauer and Bernard for a more physical interpretation.

This paper is organized as follows. First, we review several examples of commutation relations arising from different properties (general or particular) of SLE. We then consider commutation at the level of infinitesimal generators. Some (rational) solutions are discussed, and these necessary infinitesimal conditions are recast as an integrability problem. Study of integrability conditions leads to the classification of some “cocycles” on the configuration space. This leads to a holonomic system of PDEs (extending a natural situation in critical percolation), when the residual moduli space is reduced to a point. Restriction formulae are derived in a particular (chordal) case; the analogue radial case follows. Finally, one discusses corresponding questions in multiply connected domains.

1 Introduction and notations

First we recall some definitions and fix notations. We shall be mainly interested in two kinds of SLE: chordal SLE in the upper half-plane $\mathbb{H}$, from a real point to ∞ ; and radial SLE in the unit disk $\mathbb{U}$, from a boundary point to 0. Corresponding SLEs in other (simply connected) domains are obtained by conformal equivalence.

For general background on SLE, see [18, 24, 15]. Also, we will use freely results on the restriction property and the “loop soup” (see [13, 17, 22]).

Consider the family of ODEs, indexed by z in $\mathbb{H}$:

$$\partial_t g_t(z) = \frac{2}{g_t(z) - W_t}$$

with initial conditions $g_0(z) = z$, where W_t is some real-valued (continuous) function. These chordal Loewner equations are defined up to explosion time τ_z (maybe infinite). Define:

$$K_t = \overline{\{z \in \mathbb{H} : \tau_z < t\}}.$$

Then $(K_t)_{t \geq 0}$ is an increasing family of compact subsets of $\overline{\mathbb{H}}$; moreover, g_t is the unique conformal equivalence $\mathbb{H} \setminus K_t \rightarrow \mathbb{H}$ such that (hydrodynamic normalization at ∞):

$$g_t(z) = z + o(1).$$

For any compact subset K of $\overline{\mathbb{H}}$ such that $\mathbb{H} \setminus K$ is simply connected, we denote by ϕ the unique conformal equivalence $\mathbb{H} \rightarrow \mathbb{H} \setminus K$ with hydrodynamic normalization at ∞ ; so that $g_t = \phi_{K_t}$.

The coefficient of $1/z$ in the Laurent expansion of g_t at ∞ is by definition the half-plane capacity of K_t at infinity; this capacity equals $(2t)$.

If $W_t = x + \sqrt{\kappa}B_t$ where (B_t) is a standard Brownian motion, then the Loewner chain (K_t) (or the family (g_t)) defines the chordal Schramm-Loewner Evolution with parameter κ in $(\mathbb{H}, x, \infty)$. The chain K_t is generated by the trace γ , a continuous process taking values in $\overline{\mathbb{H}}$, in the following sense: $\mathbb{H} \setminus K_t$ is the unbounded connected component of $\mathbb{H} \setminus \gamma_{[0, t]}$.

The trace is a continuous non self-traversing curve. It is a.s. simple if $\kappa \leq 4$ and a.s. space-filling if $\kappa \geq 8$.

In the radial case, Loewner’s equations are indexed by $z \in \mathbb{U}$,

$$\partial_t g_t(z) = -g_t(z) \frac{g_t(z) + \xi_t}{g_t(z) - \xi_t}$$

and $g_0(z) = z$, ξ takes values in the unit circle. The hull K_t is defined as above, and g_t is the unique conformal equivalence $\mathbb{U} \setminus K_t \rightarrow \mathbb{U}$ with $g_t(0) = 0$, $g'_t(0) > 0$. Moreover, $g'_t(0) = e^{-t}$. If $\xi_t = \xi_0 \exp(i\sqrt{\kappa}B_t)$, where B is a standard Brownian motion, one gets radial SLE $_{\kappa}$ from ξ_0 to 0 in $\mathbb{U}$.

Note that chordal SLE depends only on two boundary points, and radial SLE depends on one boundary and one bulk point. In several natural instances, one needs to track additional points on the boundary. This has prompted the introduction of SLE (κ, ρ) processes in [13], generalized in [7]. The driving Brownian motion is replaced by a semimartingale which has local Girsanov density w.r.t. the original Brownian motion.

In the chordal case, let $\underline{\rho}$ be a multi-index, i.e. :

$$\underline{\rho} \in \bigcup_{i \geq 0} \mathbb{R}^i$$

Let k be the length of $\underline{\rho}$; if $k = 0$, one simply defines SLE $(\kappa, \emptyset)$ as a standard SLE $_{\kappa}$. If $k > 0$, assume the existence of processes $(W_t)_{t \geq 0}$ and $(Z_t^{(i)})_{t \geq 0}$, $i \in \{1 \dots k\}$ satisfying the SDEs:

$$\begin{cases} dW_t = \sqrt{\kappa}dB_t + \sum_{i=1}^k \frac{\rho_i}{W_t - Z_t^{(i)}} dt \\ dZ_t^{(i)} = \frac{2}{Z_t^{(i)} - W_t} dt \end{cases} \quad (1.1)$$

and such that the processes $(W_t - Z_t^{(i)})$ do not change sign. Then we define the chordal $\text{SLE}_\kappa(\underline{\rho})$ process starting from $(w, z_1, \dots, z_k)$ as a chordal Schramm-Loewner evolution the driving process of which has the same law as (W_t) as defined above, with $W_0 = w, Z_0^{(i)} = z_i$.

In the radial case, assume the existence of processes $(\xi_t)_{t \geq 0}$ and $(\chi_t^{(i)})_{t \geq 0}, i \in \{1 \dots k\}$ satisfying the SDEs:

$$\begin{cases} d\xi_t = (i\xi_t \sqrt{\kappa} dB_t - \frac{\kappa}{2} \xi_t dt) + \sum_{i=1}^k \frac{\rho_i}{2} \left(-\xi_t \frac{\xi_t + \chi_t^{(i)}}{\xi_t - \chi_t^{(i)}} \right) dt \\ d\chi_t^{(i)} = -\chi_t^{(i)} \frac{\chi_t^{(i)} + \xi_t}{\chi_t^{(i)} - \xi_t} dt \end{cases} \quad (1.2)$$

The processes $\xi, \chi^{(i)}$ may bounce on each other but not cross. This defines radial $\text{SLE}_\kappa(\underline{\rho})$ in the unit disk. Note the factor $1/2$ before the ρ_i parameters in the SDE: this is to ensure coherence with the chordal case.

2 Examples of commutation

We begin by discussing how properties of SLE (e.g. reversibility and duality) yield natural examples of commutation relations.

Reversibility: consider a chordal SLE in $(\mathbb{H}, 0, \infty)$, γ its trace. For simplicity, assume that $\kappa \leq 4$, so that the trace is a.s. simple. Define $\hat{\gamma}_t = \gamma_{1/t}$. Then $\hat{\gamma}$ is a simple curve from ∞ to 0 in $\mathbb{H}$ (for transience of SLE, see [18]). After a time change $s = s(t)$, $\hat{\gamma}$ is such that $-1/\hat{\gamma}_{[0,s]}$ has capacity $2s$. Then, according to reversibility, $\hat{\gamma}$ a SLE_κ in $\mathbb{H}$ from ∞ to 0 . (Note that for all $0 < \kappa \leq 4$ and involution of $\mathbb{H}$ of type $z \mapsto -\lambda/z$, this defines a somewhat intricate measure-preserving involution of the Wiener space).

Admitting reversibility, one can define a chordal SLE growing “from both ends” in the following fashion: let B be a standard Brownian motion, with filtration $\mathcal{F}$, γ the trace of the associated SLE_κ , and $\hat{\gamma}$ as above. Then consider $(K_{t,s} = \gamma_{[0,t]} \cup \hat{\gamma}_{[0,s]}^{-1})_{s,t \geq 0}$ and the filtration $(\mathcal{G}_{t,s})$ it generates. For any (t_0, s_0) , $(K_{t_0+t, s_0+s})_t$ is a (time-changed) chordal SLE in $\mathbb{H} \setminus K_{t_0, s_0}$, from γ_{t_0} to $\hat{\gamma}_{s_0}$; conversely $(K_{t_0, s_0+s})_s$ is a (time-changed) chordal SLE in $\mathbb{H} \setminus K_{t_0, s_0}$, from $\hat{\gamma}_{s_0}$ to γ_{t_0} .

Together with conformal equivalence, this gives the following Markov property: if $f_{t,s}$ is a conformal equivalence $(\mathbb{H} \setminus K_{t,s}, \gamma_t, \hat{\gamma}_s) \rightarrow (\mathbb{H}, 0, \infty)$ with some normalization (e.g. $f_{t,s}(1) = 1$), then $f_{t_0, s_0}(K_{t_0+t, s_0+s})$ is up to a time-change $\mathbb{R}_+^2 \rightarrow \mathbb{R}_+^2$ a copy of $(K_{t,s})$ independent from $\mathcal{G}_{t_0, s_0}$.

Duality: Duality relates the boundary of non-simple SLE ($\kappa > 4$) with corresponding simple SLEs ($\kappa' = 16/\kappa$). Let us try to formulate a precise conjecture in a “dual” fashion. We elaborate on restriction formulae identities discussed in [7].

Let $\kappa > 4, \kappa' = 16/\kappa$. Consider the configuration $(\mathbb{H}, x, y, z, \infty)$, where $x < y < z$. Define a Loewner chain from 0 to ∞ as follows:

the chain $(K_t)_{t \leq \tau_z}$ is an $\text{SLE}_\kappa(\kappa/2 - 4, -\kappa/2)$ in $\mathbb{H}$, started from (x, y, z) , aiming at ∞ , and stopped at time τ_z when the trace hits z (which it does with probability 1). Then $(K_{t+\tau_z})_{t \geq 0}$ is a $\text{SLE}_\kappa(\kappa - 4)$ in $\mathbb{H} \setminus K_{\tau_z}$, started from (z, z^+) and aiming at ∞ .

The right-boundary of $K_\infty = \bigcup_{t \geq 0} K_t$ is a simple curve from z to ∞ in $\mathbb{H}$; denote by (δ_u) the corresponding Loewner trace (i.e. $\delta_{[0, \infty)}$ is the right-boundary of K_∞ and $\delta_{[0, u]}$ has half-plane capacity $(2u)$).

Now consider a configuration $(\mathbb{H}, x', y', z', \infty)$, where $x' < y' < z'$, and let γ' be the trace of the chordal $\text{SLE}_{\kappa'}(-\kappa'/2, \kappa' - 2)$ in $\mathbb{H}$ started from (z', x', y') and aiming at ∞ .

Then we can formulate:

Conjecture 1. *The following statements hold:*

- (i) The law of δ is that of γ , where $x = x'$, $y = y'$, $z = z'$.
(ii) The law of $(\phi_{\delta_{[0,u]}}(K_t))_{t \geq 0}$ conditionally on $\delta_{[0,u]}$ is (up to a time-change) that of a copy of (K_t) started from $(x', y', z') = \phi_{\delta_{[0,u]}}(x, y, \delta_u)$.

This conjecture can be interpreted in terms of multiple SLEs: one can grow simultaneously the chain (K_t) and its (final) right-boundary. One also get a Markov property similar to the one discussed for reversibility.

Locality: The scaling limit of the exploration process for critical site percolation on the triangular lattice is SLE_6 (see [21, 4]). For some boundary conditions, one can define several exploration paths. Consider for instance the following situation: $(D, x_1, \dots, x_{2n})$ is a simply connected domain with $(2n)$ marked boundary points in cyclical order. The segments $(x_1, x_2), \dots, (x_{2n-1}, x_{2n})$ (resp. $(x_2, x_3), \dots, (x_{2n}, x_1)$) are set to blue (resp. yellow). Then one can start an exploration process at each of the boundary points x_i ; these are well-defined up to some disconnection event.

One can also consider some conditional versions: for instance, critical percolation in $(\mathbb{H}, 0, 1)$, where the half-lines $(\infty, 0)$ and $(1, \infty)$ are blue and $(0, 1)$ is yellow, conditionally on the existence of a yellow path from $(0, 1)$ to infinity (this is a singular conditioning, related to the one-arm half-plane exponent). Now the exploration processes started from 0 and 1 resp. can be defined for all time. The two traces intersect at pivotal points for the conditioning event.

One may also consider the following situation: a conformal rectangle, with sides alternately blue and yellow. Hence, one can start four exploration processes (one at each vertex). Then condition on a Cardy crossing event (e.g. the two blue sides are connected by a blue path). One can note that in this situation, the Girsanov drift terms are not rational functions.

Restriction: The restriction property of $\text{SLE}_{8/3}$ can be used to get commutation relations. For instance, consider a simply connected domain with four marked points on the boundary, say $(\mathbb{H}, a, b, c, d)$. One can define two independent $\text{SLE}_{8/3}$'s, from a to b and c to d resp., and condition them on not intersecting. Then, from the restriction property, this system of two SLEs has a natural Markov property, and also a restriction property.

More precisely, let γ and γ' be the traces of these SLEs, (g_t) the family of conformal equivalences of the first one (for some time parameterization). Then ($\mathcal{L}$ denotes probability distributions)

$$\begin{aligned} \mathcal{L}(g_t(\gamma_{(t,\infty)}), g_t(\gamma') | \gamma \cap \gamma' = \emptyset) &= \mathcal{L}(g_t(\gamma_{(t,\infty)}), g_t(\gamma') | \gamma_{(0,t)} \cap \gamma' = \emptyset, g_t(\gamma_{(t,\infty)}) \cap g_t(\gamma')) \\ &= \mathcal{L}(\tilde{\gamma}, \tilde{\gamma}' | \tilde{\gamma} \cap \tilde{\gamma}' = \emptyset) \end{aligned}$$

where $\tilde{\gamma}$ and $\tilde{\gamma}'$ are independent $\text{SLE}_{8/3}$'s going from $g_t(\gamma_t)$ to $g_t(b)$ and from $g_t(c)$ to $g_t(d)$ resp. (using the Markov property for γ and the restriction property for γ'). For the restriction property, note that, for any hull A disjoint from $\{a, b, c, d\}$:

$$\begin{aligned} \mathbb{P}((\gamma \cup \gamma')' \cap A = \emptyset | \gamma \cap \gamma' = \emptyset) &= \frac{\mathbb{P}(\gamma \cap \gamma' = \gamma \cap A = \gamma' \cap A = \emptyset)}{\mathbb{P}(\gamma \cap \gamma' = \emptyset)} \\ &= \frac{\mathbb{P}(\gamma \cap \gamma' = \emptyset | \gamma \cap A = \gamma' \cap A = \emptyset) \mathbb{P}(\gamma \cap A = \gamma' \cap A = \emptyset)}{\mathbb{P}(\gamma \cap \gamma' = \emptyset)} \\ &= \frac{\mathbb{P}(\phi_A(\gamma) \cap \phi_A(\gamma') = \emptyset) \mathbb{P}(\gamma \cap A = \gamma' \cap A = \emptyset)}{\mathbb{P}(\gamma \cap \gamma' = \emptyset)} \\ &= \mathbb{P}(\gamma \cap A = \emptyset) \mathbb{P}(\gamma' \cap A = \emptyset) \frac{\psi(\phi_A(a, b, c, d))}{\psi(a, b, c, d)} \end{aligned}$$

where $\psi(a, b, c, d)$ is the probability that the two independent $\text{SLE}_{8/3}$ do not intersect.

If $\kappa \in (0, 8/3)$, one can consider two independent SLE_κ , a corresponding independent loop soup, and condition on the event: no loop intersects the two SLEs. A standard computation shows that the probability

ψ_κ of this event is given by:

$$\psi_\kappa(\infty, 0, x, 1) = \frac{\Gamma(4/\kappa)\Gamma(12/\kappa - 1)}{\Gamma(8/\kappa)\Gamma(8/\kappa - 1)} x^{2/\kappa} {}_2F_1\left(\frac{4}{\kappa}, 1 - \frac{4}{\kappa}; \frac{8}{\kappa}; x\right)$$

In a domain with $(2n)$ marked points on the boundary in cyclical order, say $(\mathbb{H}, x_1, \dots, x_{2n})$, for a given pairing of $\{x_1, \dots, x_{2n}\}$, define n independent SLE_κ , with endpoints determined by the pairing. Consider $(n-1)$ auxiliary independent loop $L_1, \dots, L_{n-1}$ soups, with intensity λ_κ . One can consider the event: for $k = 1, \dots, n-1$, no loop in L_k intersects more than k SLEs. This has positive probability iff the pairing is a non-crossing one. As is well known, there are C_n of these pairings, where C_n is the n -th Catalan's number:

$$C_n = \frac{\binom{2n}{n}}{n+1}$$

Then one can condition on this event to get n non intersecting SLE_κ 's, that have an appropriate Markov property and restriction property. This situation is discussed in details in [10]; when $\kappa = 2$, this is directly connected to Fomin's formulae [11, 12].

Wilson's algorithm: In the case $\kappa = 8$, $\kappa' = 2$, the Uniform Spanning Tree and the Loop-Erased Random Walk converge to SLE_8 and SLE_2 resp. (see [16]). As pointed out in [16], duality follows from these convergence and Wilson's algorithm, that gives an exact relation between UST and LERW at the discrete level ([25]).

Let us formulate a precise duality identity in this situation. Consider (K_t) a chordal SLE_8 in $(\mathbb{H}, 0, \infty)$. Let G be the (random) leftmost point visited by this SLE before τ_1 . Then a standard SLE computation (see e.g. [24]) yields:

$$\mathbb{P}(G \leq -g) = \frac{1}{\pi} \int_0^g \frac{dt}{(1+t)\sqrt{t}} = \frac{2}{\pi} \arctan(\sqrt{g})$$

This distribution is the exit distribution of a random walk with normal reflection on $\mathbb{R}^+$, absorbed on $\mathbb{R}^-$, and started from 1 (as is readily seen by mapping $\mathbb{H}$ to a quadrant by $z \mapsto \sqrt{z}$ and a reflection argument).

At the discrete level, we are considering a UST wired on $\mathbb{R}^-$ and free on $\mathbb{R}^+$. The branch connecting 1 to $\mathbb{R}^-$ is a LERW started from 1 and reflected on $\mathbb{R}^+$. By a slight modification of the arguments of [16] (considering the Poisson kernel for this reflected random walk gives "harmonic martingales" for the time-reverted LERW), one gets that conditionally on G , the boundary of K_{τ_1} , which is a random simple curve connecting G and 1, and the scaling limit of this LERW, is chordal $\text{SLE}_2(-1, -1)$ in $(\mathbb{H}, G, 1)$ started from $(G, 0, \infty)$.

Wilson's algorithm gives more information. The boundary ∂K_{τ_1} divides $\mathbb{H}$ in two simply connected domains $\mathbb{H}_0$ and $\mathbb{H}_\infty$, with 0 and ∞ in their respective boundary. Then, conditionally on ∂K_{τ_1} , the original SLE_8 is the concatenation of a chordal SLE_8 in $(\mathbb{H}_0, 0, 1)$ and a chordal SLE_8 in $(\mathbb{H}_\infty, 1, \infty)$.

For small times, the law of the original SLE_8 conditionally on $G = g$ (in the regular conditional probability sense) is easy to work out. Consider the following martingale (with usual notations):

$$\arctan \sqrt{\frac{W_t - g_t(x)}{g_t(1) - W_t}}$$

Differentiating w.r.t x , one gets a local martingale:

$$\frac{g'_t(x)}{g_t(1) - g_t(x)} \sqrt{\frac{W_t - g_t(x)}{g_t(1) - W_t}}$$

Using this as a Girsanov density, one finds that the conditional SLE_8 is a chordal $\text{SLE}_8(-4, 4)$ in $(\mathbb{H}, 0, \infty)$ started from $(0, G, 1)$.

For symmetry, and from reversibility, the chordal SLE₈ in $(\mathbb{H}_\infty, 1, \infty)$ is a time-reversed chordal SLE₈ in $(\mathbb{H}_\infty, \infty, 1)$.

Consider now these different processes as chordal SLEs in $\mathbb{H}$ aiming at 1 (and not ∞). Then we have
 an SLE₂(−1, −1) started from $(G, 0, \infty)$
 an SLE₈(−4, 2) started from $(0, G, \infty)$
 an SLE₈(−4, 2) started from $(\infty, G, 0)$
 We shall see later that this fits in infinitesimal relations for SLE_κ(ρ).

One can extend the situation as follows: in the discrete setting, consider n points on $\mathbb{R}^+$ and n points on $\mathbb{R}^-$:

$$x_n < \dots < x_1 < 0 < y_1 < \dots < y_n.$$

Consider a UST with the same boundary conditions as before, and the smallest subtree containing $y_1, \dots, y_n$ and $\mathbb{R}^-$. Condition on the event that this subtree has no triple point in the bulk. Then it consists in the union of n disjoint paths in the bulk and $\mathbb{R}^-$. Now condition on the endpoints of these branches being $x_n, \dots, x_1$, and take this to the scaling limit. Using Wilson's algorithm and Fomin's formulae ([11]), everything can be made explicit, and this defines n “non-intersecting” SLE₂'s in the upper half-plane (with $(2n + 2)$ marked points on the boundary).

3 Commutation of infinitesimal generators

3.1 The commutation framework

We have seen natural examples where two SLEs could be grown in a common domain in a consistent fashion. In this section, we discuss necessary infinitesimal conditions. We shall define a “global” commutation condition, of geometric nature, and express its consequence in terms of infinitesimal generators, which is of algebraic nature.

Let us consider the following chordal situation: the domain is $\mathbb{H}$, SLEs aim at ∞ , and $(x, y, z_1, \dots, z_n)$ are $(n + 2)$ (distinct) points on the real line; the point at infinity is also a marked point. We want to grow two infinitesimal hulls (with capacity of order ε) at x and z respectively. We can either grow a hull K_ε at x , and then another one at y in the perturbed domain $\mathbb{H} \setminus K_\varepsilon$, or proceed in the other order. The coherence condition is that these two procedures yield the same result.

Let us make things more rigorous. Consider a Loewner chain $(K_{s,t})_{(s,t) \in \mathcal{T}}$ with a double time index, so that $K_{s,t} \subset K_{s',t'}$ if $s' \geq s, t' \geq t$ and $K_{s,t} \neq K_{s',t'}$ if $(s', t') \neq (s, t)$. We only consider chains up to time reparameterization $\mathbb{R}_+^2 \rightarrow \mathbb{R}_+^2$. We also assume that $K_{s,t} = K_{s,0} \cup K_{0,t}$. The time set $\mathcal{T}$ may be random, but includes a.s. a neighbourhood of $(0, 0)$ in $\mathbb{R}_+^2$. Also, if $s \leq s', t \leq t', (s', t') \in \mathcal{T}$, then $(s, t) \in \mathcal{T}$. Define $g_{s,t}$ the conformal equivalence $\mathbb{H} \setminus K_{s,t} \rightarrow \mathbb{H}$ with hydrodynamic normalization at infinity ($g_{s,t} = \phi_{K_{s,t}}$ with the earlier notation), and the continuous traces $\gamma, \tilde{\gamma}$, such that:

$$\gamma_{s,t} = \lim_{\varepsilon \searrow 0} \overline{g_{s,t}(K_{s+\varepsilon,t} \setminus K_{s,t})}, \tilde{\gamma}_{s,t} = \lim_{\varepsilon \searrow 0} \overline{g_{s,t}(K_{s,t+\varepsilon} \setminus K_{s,t})}$$

where $\gamma_{0,t} = x$ for all $(0, t) \in \mathcal{T}$, and similarly $\tilde{\gamma}_{s,0} = y$ for all $(s, 0) \in \mathcal{T}$.

Furthermore, assume that the following conditions are satisfied:

Definition 2. Let $(K_{s,t})_{(s,t) \in \mathcal{T}}$ be a random Loewner chain with double time indexing; the associated conformal equivalences are $g_{s,t} = \phi_{K_{s,t}}$. We say that $(K_{s,t})$ is an SLE($\kappa, b, \tilde{\kappa}, \tilde{b}$) if:

- (i). The time set $\mathcal{T}$ is a.s. open, connected, and a neighbourhood of $(0, 0)$ in $\mathbb{R}_+^2$. The ranges of the traces $\gamma_{\mathcal{T}}, \tilde{\gamma}_{\mathcal{T}}$ are disjoint and $z_1, \dots, z_n \notin K_{s,t}$ for $(s, t) \in \mathcal{T}$.

(ii). Let σ (resp. τ) be a stopping time in the filtration generated by $(K_{s,0})_{(s,0) \in \mathcal{T}}$ (resp. $(K_{0,t})_{(0,t) \in \mathcal{T}}$). Let also $\mathcal{T}' = \{(s,t) : (s+\sigma, t+\tau) \in \mathcal{T}\}$ and $(K'_{s,t})_{(s,t) \in \mathcal{T}'} = \left(\overline{g_{\sigma,\tau}(K_{s+\sigma, t+\tau} \setminus K_{s,t})} \right)$. Then $(K'_{s,0})_{(s,0) \in \mathcal{T}'}$ is distributed as a stopped $\text{SLE}_\kappa(b)$, i.e an SLE driven by:

$$dX_s = \sqrt{\kappa} dB_s + b(X_s, g_s(y), \dots, g_s(z_i), \dots) dt$$

Likewise $(K'_{0,t})_{(0,t) \in \mathcal{T}'}$ is distributed as a stopped $\text{SLE}_{\tilde{\kappa}}(\tilde{b})$, i.e an SLE driven by:

$$dY_t = \sqrt{\tilde{\kappa}} d\tilde{B}_t + \tilde{b}(\tilde{g}_t(x), Y_t, \dots, \tilde{g}_t(z_i), \dots) dt$$

Here $B, \tilde{B}$ are standard Brownian motions, $(g_s), (\tilde{g}_t)$ are the associated conformal equivalences, $b, \tilde{b}$ are some smooth, translation invariant, and homogeneous of degree (-1) functions. If A^x, A^y are two increasing functions of hulls growing at x and y resp. (e.g. the half-plane capacity), we shall be particularly interested in stopping times of type $\sigma = \inf(s : A^x(K_{s,0}) \geq a^x)$, $\tau = \inf(t : A^y(K_{0,t}) \geq a^y)$.

Note that $(X_s, \dots, g_t(z_i), \dots)$ is a Markov process. Let P be its semigroup and $\mathcal{L}$ its infinitesimal generator. Similarly, $(\tilde{g}_t(\hat{x}), Y_t, \dots)$ is a Markov process with semigroup Q and infinitesimal generator $\mathcal{M}$. We are interested in what conditions on the functions b and $\tilde{b}$ are implied by these assumptions (the existence of an $\text{SLE}(\kappa, b, \tilde{\kappa}, \tilde{b})$).

So let F be a test function $\mathbb{R}^{n+2} \rightarrow \mathbb{R}$, and $c > 0$ be some constant (ratio of speeds). We apply the previous assumptions with $A^x = A^y = \text{cap}$ (the half-plane capacity), $a^x = 2\varepsilon$, $a^y = 2c\varepsilon$. We are interested in the hull $K_{\sigma,\tau}$. Two ways of getting from $K_{0,0}$ to $K_{\sigma,\tau}$ are (symbolically):

$$\begin{array}{ccc} K_{0,0} & \longrightarrow & K_{\sigma,0} \\ \downarrow & & \downarrow \\ K_{0,\tau} & \longrightarrow & K_{\sigma,\tau} \end{array}$$

and our assumptions give a description of these transitions.

So consider the following procedure:

- run the first SLE (i.e. $\text{SLE}_\kappa(b)$), started from $(x, y, \dots, z_i, \dots)$ until it reaches capacity 2ε .
- then run independently the second SLE (i.e. $\text{SLE}_{\tilde{\kappa}}(\tilde{b})$) in $g_\varepsilon^{-1}(\mathbb{H})$ until it reaches capacity $2c\varepsilon$; this capacity is measured *in the original half-plane*. Let $\tilde{g}_\varepsilon$ be the corresponding conformal equivalence.
- one gets two hulls resp. at x and y with capacity 2ε and $2c\varepsilon$; let $\phi = \tilde{g}_\varepsilon \circ g_\varepsilon$ be the normalized map removing these two hulls.
- expand $\mathbb{E}(F(\tilde{g}_\varepsilon(X_\varepsilon), \tilde{Y}_\varepsilon))$ up to order two in ε .

This describes (in distribution) how to get from $K_{0,0}$ to $K_{\sigma,0}$, and then from $K_{\sigma,0}$ to $K_{\sigma,\tau}$.

From the Loewner equation, it appears that $\partial_t g'_t(w) = -2g'_t(w)/(g_t(w) - W_t)^2$. Hence $g'_\varepsilon(y) = 1 - 2\varepsilon/(y - x)^2 + o(\varepsilon)$. From the scaling property of half-plane capacity, we get:

$$\tilde{\varepsilon} = c\varepsilon \left(1 - \frac{4\varepsilon}{(y-x)^2} \right) + o(\varepsilon^2)$$

i.e $\tilde{\varepsilon}$ is deterministic up to order two in ε . Denote by $\mathcal{L}$ and $\mathcal{M}$ the infinitesimal generators of the two SLEs:

$$\begin{aligned} \mathcal{L} &= \frac{\kappa}{2} \partial_{xx} + b(x, y, \dots) \partial_x + \frac{2}{y-x} \partial_y + \sum_{i=1}^n \frac{2}{z_i - x} \partial_i \\ \mathcal{M} &= \frac{\tilde{\kappa}}{2} \partial_{yy} + \tilde{b}(x, y, \dots) \partial_y + \frac{2}{x-y} \partial_x + \sum_{i=1}^n \frac{2}{z_i - y} \partial_i \end{aligned}$$

where $\partial_i = \partial_{z_i}$. Let $w = (x, y, \dots, z_i, \dots)$, $w' = (X_\varepsilon, g_\varepsilon(y), \dots, g_\varepsilon(z_i), \dots)$, $w'' = (\tilde{g}_\varepsilon(X_\varepsilon), \tilde{Y}_\varepsilon, \dots, \tilde{g}_\varepsilon \circ g_\varepsilon(z_i), \dots)$. Now:

$$\begin{aligned} \mathbb{E}(F(w'')|w) &= \mathbb{E}(F(w'')|w'|w) = P_\varepsilon \mathbb{E}(Q_\varepsilon F|w')(w) \\ &= P_\varepsilon \mathbb{E} \left(\left(1 + \tilde{\varepsilon} \mathcal{M} + \frac{\tilde{\varepsilon}^2}{2} \mathcal{M}^2\right) F(w') \right) (w) = P_\varepsilon Q_{c\varepsilon(1-4\varepsilon/(y-x)^2)} F(w) + o(\varepsilon^2) \\ &= \left(1 + \varepsilon \mathcal{L} + \frac{\varepsilon^2}{2} \mathcal{L}^2\right) \left(1 + c\varepsilon \left(1 - \frac{4\varepsilon}{(y-x)^2}\right) \mathcal{M} + \frac{c^2 \varepsilon^2}{2} \mathcal{M}^2\right) F(w) + o(\varepsilon^2) \\ &= \left(1 + \varepsilon(\mathcal{L} + c\mathcal{M}) + \varepsilon^2 \left(\frac{1}{2} \mathcal{L}^2 + \frac{c^2}{2} \mathcal{M}^2 + c\mathcal{L}\mathcal{M} - \frac{4c}{(y-x)^2} \mathcal{M}\right)\right) F(w) + o(\varepsilon^2) \end{aligned}$$

If we first grow a hull at z , then at x , one gets instead:

$$\left(1 + \varepsilon(\mathcal{L} + c\mathcal{M}) + \varepsilon^2 \left(\frac{1}{2} \mathcal{L}^2 + \frac{c^2}{2} \mathcal{M}^2 + c\mathcal{M}\mathcal{L} - \frac{4c}{(z-x)^2} \mathcal{L}\right)\right) F(w) + o(\varepsilon^2)$$

Hence the commutation condition reads:

$$[\mathcal{L}, \mathcal{M}] = \frac{4}{(y-x)^2} (\mathcal{M} - \mathcal{L})$$

. After simplifications, one gets:

$$\begin{aligned} [\mathcal{L}, \mathcal{M}] + \frac{4}{(y-x)^2} (\mathcal{L} - \mathcal{M}) &= (\kappa \partial_x \tilde{b} - \tilde{\kappa} \partial_y b) \partial_{xy} \\ &+ \left[\frac{2\partial_x b}{y-x} + \sum_i \frac{2\partial_i b}{y-z_i} - \tilde{b} \partial_y b + \frac{2b}{(y-x)^2} + \frac{2\kappa-12}{(x-y)^3} - \frac{\tilde{\kappa}}{2} \partial_{yy} b \right] \partial_x \\ &- \left[\frac{2\partial_y \tilde{b}}{x-y} + \sum_i \frac{2\partial_i \tilde{b}}{x-z_i} - b \partial_x \tilde{b} + \frac{2\tilde{b}}{(x-y)^2} + \frac{2\tilde{\kappa}-12}{(y-x)^3} - \frac{\kappa}{2} \partial_{xx} \tilde{b} \right] \partial_y \end{aligned}$$

So the commutation condition reduces to three differential conditions involving b and $\tilde{b}$; note the non-linear terms $\tilde{b} \partial_y b$ and $b \partial_x \tilde{b}$.

3.2 Rational solutions

Case $n = 0$:

If $n = 0$, then $b(x, y) = \rho/(x-y)$ and $\tilde{b}(x, y) = \tilde{\rho}/(y-x)$. Then the commutation condition reduces to:

$$\begin{cases} \kappa \tilde{\rho} = \tilde{\kappa} \rho \\ (\tilde{\kappa} - 4)\rho - \rho \tilde{\rho} + 12 - 2\kappa = 0 \\ (\kappa - 4)\tilde{\rho} - \rho \tilde{\rho} + 12 - 2\tilde{\kappa} = 0 \end{cases}$$

We are only interested in the case $\kappa, \tilde{\kappa} > 0$. Then:

$$\begin{cases} \tilde{\rho} = \tilde{\kappa} \rho / \kappa \\ (\tilde{\kappa} - 4)\rho - \rho^2 \tilde{\kappa} / \kappa + 12 - 2\kappa = 0 \\ (\kappa - 4)\rho \tilde{\kappa} / \kappa - \rho^2 \tilde{\kappa} / \kappa + 12 - 2\tilde{\kappa} = 0 \end{cases}$$

The last two are polynomials in ρ that have a common root if and only if their resultant vanishes. This resultant (a polynomial in the coefficients) equals:

$$\frac{12\tilde{\kappa}(\tilde{\kappa} - \kappa)^2(\kappa\tilde{\kappa} - 16)}{\kappa^3}$$

So either $\tilde{\kappa} = \kappa$, and then $\rho = \tilde{\rho} \in \{2, \kappa - 6\}$, or $\tilde{\kappa} = 16/\kappa$, and then $\rho = -\kappa/2$, $\tilde{\rho} = -\tilde{\kappa}/2$.

Let us comment briefly on these solutions. The condition $\kappa\tilde{\kappa} = 16$ obviously points at duality. The case $\tilde{\kappa} = \kappa$, $\tilde{\rho} = \rho = \kappa - 6$ corresponds in fact to reversibility. Indeed, one has:

Lemma 3. *An $\text{SLE}_\kappa(\rho_1, \dots, \rho_n, \rho)$ in $\mathbb{H}$ started from $(x, z_1, \dots, z_n, y)$ and aiming at ∞ is identical in law to a (time-changed) $\text{SLE}_\kappa(\rho_1, \dots, \rho_n, \kappa - 6 - \rho - \sum_i \rho_i)$ in $\mathbb{H}$ started from $(x, z_1, \dots, z_n, \infty)$ and aiming at y , up to disconnection of y .*

Proof. Let (g_t) be the family of conformal equivalences for the first SLE, (K_t) the corresponding hulls, W its driving process, $Y_t^{(i)} = g_t(z_i)$, $Y_t = g_t(y)$. Consider the homographies:

$$\varphi_t(u) = \frac{g'_t(y)}{Y_t - u} - \frac{g''_t(y)}{2g'_t(y)}$$

and $\tilde{g}_t = \phi_{\varphi_0(K_t)}$. Then $\tilde{g}_t \circ \varphi_0 = \varphi_t \circ g_t$, and $(\tilde{g}_t)$ defines a time-changed Loewner chain. Let $\tilde{W}_t = \varphi_t(W_t)$, $\tilde{Y}_t^{(i)} = \varphi_t(Y_t^{(i)})$, $V_t = \varphi_t(\infty)$. Then, from Itô's formula:

$$\begin{aligned} d\tilde{W}_t &= d\left(\frac{g'_t(y)}{Y_t - W_t} - \frac{g''_t(y)}{2g'_t(y)}\right) = \frac{g'_t(y)}{(Y_t - W_t)} \left(-\frac{2dt}{(Y_t - W_t)^2} - \frac{d(Y_t - W_t)}{Y_t - W_t} + \frac{\kappa dt}{(Y_t - W_t)^2} - \frac{2dt}{(Y_t - W_t)^2}\right) \\ &= \frac{g'_t(y)}{(Y_t - W_t)} \left(\frac{dW_t}{Y_t - W_t} + \frac{(\kappa - 6)dt}{(Y_t - W_t)^2}\right) \\ &= \frac{g'_t(y)\sqrt{\kappa}}{(Y_t - W_t)^2} dB_t + \frac{g'_t(y)^2}{(Y_t - W_t)^4} \left(\sum_i \rho_i \frac{Y_t - W_t}{Y_t^{(i)} - W_t} \cdot \frac{1}{\tilde{W}_t - V_t} + \frac{\kappa - 6 - \rho}{\tilde{W}_t - V_t}\right) dt \end{aligned}$$

Note that the cross-ratio is conformally invariant, so

$$\frac{Y_t - W_t}{Y_t^{(i)} - W_t} = [\infty, Y_t^{(i)}, Y_t, W_t] = [V_t, \tilde{Y}_t^{(i)}, \infty, \tilde{W}_t] = \frac{\tilde{Y}_t^{(i)} - V_t}{\tilde{Y}_t^{(i)} - \tilde{W}_t}$$

After a time change $ds = g'_t(y)^2 dt / (Y_t - W_t)^4$, one gets:

$$\begin{aligned} d\tilde{W}_s &= \sqrt{\kappa} d\tilde{B}_s + \sum_i \rho_i \frac{\tilde{Y}_t^{(i)} - V_t}{(\tilde{Y}_t^{(i)} - \tilde{W}_t)(\tilde{W}_t - V_t)} dt + \frac{\kappa - 6 - \rho}{\tilde{W}_t - V_t} dt \\ &= \sqrt{\kappa} d\tilde{B}_s + \sum_i \frac{\rho_i}{\tilde{Y}_t^{(i)} - \tilde{W}_t} dt + \frac{\kappa - 6 - \rho - \sum_i \rho_i}{\tilde{W}_t - V_t} dt \end{aligned}$$

which defines a $\text{SLE}_\kappa(\rho_1, \dots, \rho_n, \kappa - 6 - \rho - \sum_i \rho_i)$. □

In particular, an $\text{SLE}_\kappa(\kappa - 6)$ started from (x, y) and stopped at τ_y is simply an SLE_κ from x to y . For $\kappa = 6$ and $n = 0$, this is locality.

Parametric case:

Assume the following forms for the drift terms $b, \tilde{b}$:

$$\begin{aligned} b(x, y, z_i) &= \frac{\rho}{x - y} + \sum_i \frac{\rho_i}{x - z_i} \\ \tilde{b}(x, y, z_i) &= \frac{\tilde{\rho}}{y - x} + \sum_i \frac{\tilde{\rho}_i}{y - z_i} \end{aligned}$$

Then the commutation conditions are:

$$\begin{cases} \kappa\tilde{\rho} = \tilde{\kappa}\rho \\ (\tilde{\kappa} - 4)\rho - \rho\tilde{\rho} + 12 - 2\kappa = 0 \\ (\kappa - 4)\tilde{\rho} - \rho\tilde{\rho} + 12 - 2\tilde{\kappa} = 0 \\ \rho\tilde{\rho}_i = 2\rho_i \\ \tilde{\rho}\rho_i = 2\tilde{\rho}_i \end{cases}$$

As we have seen, the first three conditions imply that $\kappa = \tilde{\kappa}$, $\rho = \tilde{\rho} \in \{2, \kappa - 6\}$, or $\kappa\tilde{\kappa} = 16$, $\rho = -\kappa/2$, $\tilde{\rho} = -\tilde{\kappa}/2$. Now, if the $\rho_i, \tilde{\rho}_i$ are not all zero, $\rho\tilde{\rho} = 4$, which happens if $\rho = \tilde{\rho} = 2$, or in the case $\kappa\tilde{\kappa} = 16$. To sum up, the solutions are:

- (i) $\kappa = \tilde{\kappa}$, $\rho = \tilde{\rho} = \kappa - 6$, $\rho_i = \tilde{\rho}_i = 0$ (1 free parameter)
- (ii) $\kappa = \tilde{\kappa}$, $\rho = \tilde{\rho} = 2$, $\rho_i = \tilde{\rho}_i$ $((1+n)$ free parameters)
- (iii) $\kappa\tilde{\kappa} = 16$, $\rho = -\kappa/2$, $\tilde{\rho} = -\tilde{\kappa}/2$, $\tilde{\rho}_i = -(\tilde{\kappa}/4)\rho_i = -(4/\kappa)\rho_i$ $((1+n)$ free parameters)

These examples are “rational” (i.e. the drift terms are rational functions). Yet, important examples (deduced from locality and restriction) are transcendental. In the next section, we recast these commutation conditions as integrability conditions, satisfied by all these examples.

4 Integrability conditions

4.1 Integrability for SLE commutation relations

In the previous paragraph, we derived the following commutation conditions:

$$\begin{cases} \kappa\partial_x\tilde{b} - \tilde{\kappa}\partial_y b = 0 \\ \frac{2\partial_x b}{y-x} + \sum_i \frac{2\partial_i b}{y-z_i} - \tilde{b}\partial_y b + \frac{2b}{(y-x)^2} + \frac{2\kappa-12}{(x-y)^3} - \frac{\tilde{\kappa}}{2}\partial_{yy}b = 0 \\ \frac{2\partial_y\tilde{b}}{x-y} + \sum_i \frac{2\partial_i\tilde{b}}{x-z_i} - b\partial_x\tilde{b} + \frac{2\tilde{b}}{(x-y)^2} + \frac{2\tilde{\kappa}-12}{(y-x)^3} - \frac{\kappa}{2}\partial_{xx}\tilde{b} = 0 \end{cases}$$

Now, from the first equation, one can write:

$$b = \kappa \frac{\partial_x \psi}{\psi}, \tilde{b} = \tilde{\kappa} \frac{\partial_y \psi}{\psi}$$

for some non-vanishing function ψ (at least locally).

It turns out that the second condition now writes:

$$-\kappa\partial_x \left(\frac{\frac{\tilde{\kappa}}{2}\partial_{yy}\psi + \sum_i \frac{2\partial_i\psi}{z_i-y} + \frac{2\partial_x\psi}{x-y} + (1-\frac{6}{\kappa})\frac{\psi}{(x-y)^2}}{\psi} \right) = 0.$$

Symmetrically, the last equation is:

$$-\tilde{\kappa}\partial_y \left(\frac{\frac{\kappa}{2}\partial_{xx}\psi + \sum_i \frac{2\partial_i\psi}{z_i-x} + \frac{2\partial_y\psi}{y-x} + (1-\frac{6}{\tilde{\kappa}})\frac{\psi}{(y-x)^2}}{\psi} \right) = 0.$$

This means that a non-vanishing solution of

$$\begin{cases} \frac{\kappa}{2}\partial_{xx}\psi + \sum_i \frac{2\partial_i\psi}{z_i-x} + \frac{2\partial_y\psi}{y-x} + \left(\left(1-\frac{6}{\kappa}\right) \frac{1}{(y-x)^2} + h_1(x, z) \right) \psi = 0 \\ \frac{\tilde{\kappa}}{2}\partial_{yy}\psi + \sum_i \frac{2\partial_i\psi}{z_i-y} + \frac{2\partial_x\psi}{x-y} + \left(\left(1-\frac{6}{\tilde{\kappa}}\right) \frac{1}{(x-y)^2} + h_2(y, z) \right) \psi = 0 \end{cases}$$

yields drift terms $b, \tilde{b}$ that satisfy the commutation condition. Obviously, these differential operators are infinitesimal generators of the SLEs, with an added coefficient before the constant term.

The problem is now to find functions h_1, h_2 such that the above system has solutions (integrability conditions). Note that we have not considered yet the conditions: $b, \tilde{b}$ translation invariant and homogeneous of degree (-1) . This implies that ψ can be chosen to be translation invariant and homogeneous of some fixed degree. So assume that we are given h_1, h_2 , and a non-vanishing (translation-invariant, homogeneous) solution ψ of this system. Let:

$$\mathcal{M}_1 = \frac{\kappa}{2} \partial_{xx} \psi + \sum_i \frac{2\partial_i \psi}{z_i - x} + \frac{2\partial_y \psi}{y - x} + \left(1 - \frac{6}{\tilde{\kappa}}\right) \frac{1}{(y - x)^2}, \mathcal{M}_2 = \frac{\tilde{\kappa}}{2} \partial_{yy} \psi + \sum_i \frac{2\partial_i \psi}{z_i - y} + \frac{2\partial_x \psi}{x - y} + \left(1 - \frac{6}{\kappa}\right) \frac{1}{(x - y)^2}$$

Then ψ is annihilated by all operators in the left ideal generated by $(\mathcal{M}_1 + h_1), (\mathcal{M}_2 + h_2)$, including in particular:

$$\begin{aligned} \mathcal{M} &= [\mathcal{M}_1 + h_1, \mathcal{M}_2 + h_2] + \frac{4}{(x - y)^2} ((\mathcal{M}_1 + h_1) - (\mathcal{M}_2 + h_2)) \\ &= [\mathcal{M}_1, \mathcal{M}_2] + \frac{4}{(x - y)^2} (\mathcal{M}_1 - \mathcal{M}_2) + ([\mathcal{M}_1, h_2] - [\mathcal{M}_2, h_1]) + \frac{4(h_1 - h_2)}{(x - y)^2} \\ &= \frac{3(\kappa\tilde{\kappa} - 16)(\kappa - \tilde{\kappa})}{\kappa\tilde{\kappa}(x - y)^4} + \left(\left(\frac{2\partial_y}{y - x} + \sum_i \frac{2\partial_i}{z_i - x} \right) h_2 - \left(\frac{2\partial_x}{x - y} + \sum_i \frac{2\partial_i}{z_i - y} \right) h_1 \right) + \frac{4(h_1 - h_2)}{(x - y)^2} \end{aligned}$$

This is an operator of order 0, so it must vanish identically. Considering the pole at $x = y$, this implies in particular $\tilde{\kappa} \in \{\kappa, 16/\kappa\}$, since the fourth-order pole must vanish. Then the second-order pole must also vanish, so $h_1(x, z) = h(x, z)$, $h_2(y, z) = h(y, z)$ for some h . So this condition boils down to a functional equation on h .

For illustration, consider the following variation on an earlier example: a chordal SLE_{8/3} from x to y is conditioned not to intersect an independent restriction measure from z to ∞ with index ν . Let $\varphi(x, y, z)$ be the probability of non-intersection. Then φ is annihilated by the operator:

$$\frac{\kappa}{2} \partial_{xx} + \frac{\kappa - 6}{x - y} \partial_x + \frac{2}{y - x} \partial_y + \frac{2}{z - x} \partial_z - \frac{2\nu}{(x - z)^2}$$

where $\kappa = 8/3$. Obviously φ can be expressed in terms of a hypergeometric function. If $\psi = (y - x)^{-2\alpha} \varphi$, $\alpha = \alpha_\kappa = 5/8$, then ψ is annihilated by the conjugate operators:

$$\frac{\kappa}{2} \partial_{xx} + \frac{2}{y - x} \partial_y + \frac{2}{z - x} \partial_z - \frac{2\alpha}{(x - y)^2} - \frac{2\nu}{(x - z)^2}, \frac{\kappa}{2} \partial_{yy} + \frac{2}{x - y} \partial_x + \frac{2}{z - y} \partial_z - \frac{2\alpha}{(y - x)^2} - \frac{2\nu}{(y - z)^2}$$

where we also use reversibility for SLE_{8/3}. It is easy to check that in general $\tilde{\kappa} \in \{\kappa, 16/\kappa\}$, $h(x, z) = -2\nu/(x - z)^2$ is a solution of the integrability condition above. More generally, if n points $z_1, \dots, z_n$ are marked on the real line, a (particular) solution of the integrability condition is given by $\tilde{\kappa} \in \{\kappa, 16/\kappa\}$,

$$h(x, z) = -2 \sum_i \frac{\mu_i}{(z_i - x)^2} - 2 \sum_{i < j} \nu_{ij} \left(\frac{1}{z_i - x} - \frac{1}{z_j - x} \right)^2 = \sum_i \frac{\mu'_i}{(z_i - x)^2} + \sum_{i < j} \frac{\nu'_{ij}}{(z_i - x)(z_j - x)}$$

where μ_i, ν_{ij} are real parameters. When $\kappa = \tilde{\kappa} = 8/3$, $\mu_i, \nu_{ij} \geq 0$, and $x < y < z_1 < \dots < z_n$, it is easy to think of a probabilistic situation corresponding to this. Consider a chordal SLE_{8/3} from x to y , and condition it not to intersect independent one-sided restriction samples $z_i \leftrightarrow \infty$ (with index μ_i) and $z_i \leftrightarrow z_j$ (with index $\nu_{i,j}$). Then reversibility for the conditional SLE corresponds to a partition function ψ solving PDEs where h is as above.

Let us get back to the functional equation for h :

$$\left(\left(\frac{\partial_y}{y-x} + \sum_i \frac{\partial_i}{z_i-x} \right) h(y, z) - \left(\frac{\partial_x}{x-y} + \sum_i \frac{\partial_i}{z_i-y} \right) h(x, z) \right) + 2 \frac{h(x, z) - h(y, z)}{(x-y)^2} = 0. \quad (4.3)$$

We want to prove that the only solutions to this functional equation (translation invariant and homogeneous of degree (-2)) are the rational functions given above when there are at most 3 marked z points (including infinity). By expanding in ε where $y = x + \varepsilon$, one sees that h must be annihilated by the family of operators:

$$\ell_{0,n} = \frac{n}{(n+1)(n+2)} \partial_x^{n+2} + \sum_i \left(\frac{\partial_i \partial_x^n}{z_i - x} - \frac{n! \partial_i}{(z_i - x)^{n+1}} \right)$$

for $n \geq 0$. Also, h must be translation invariant and homogeneous of degree -2 . So for $n = 2$, one can write $h(x, y, z) = \tilde{h}((y-x)/(z-x))/(z-x)^2$ for instance, and $\tilde{h}$ satisfies a third-order ODE (since $\ell_{0,1}h = 0$); but we have already exhibited 3 linearly independent solutions in this case, so the classification is complete for $n = 1, 2$. In particular, in the case $n = 2$, this is closely related to the discussion in Section 8.5 of [13].

When $n \geq 3$, the configuration $(\mathbb{H}, z_1, \dots, z_n, \infty)$ corresponds to a $(n-2)$ -dimensional moduli space. We already know $n(n+1)/2$ (linearly independent) rational solutions. We proceed to show that arbitrary (smooth) functions on this “residual moduli space” lead to solutions of the functional equation.

Define ℓ_x, ℓ_y to be the differential operators:

$$\ell_x = \frac{2}{y-x} \partial_y + \sum_i \frac{2}{z_i-x} \partial_i, \quad \ell_y = \frac{2}{x-y} \partial_x + \sum_i \frac{2}{z_i-y} \partial_i$$

representing the Loewner flow with singularities at x, y respectively, restricted to marked points. Now the operators

$$\tilde{\ell}_x = \frac{6}{y-x} \partial_x + \ell_x, \quad \tilde{\ell}_y = \frac{6}{x-y} \partial_y + \ell_y$$

are the generators of $\text{SLE}_0(-6)$, i.e. the hyperbolic geodesic from x to y . In this case, we have seen that the commutation relation:

$$[\tilde{\ell}_x, \tilde{\ell}_y] = \frac{4}{(x-y)^2} (\tilde{\ell}_y - \tilde{\ell}_x)$$

is satisfied. Hence if f is a positive, translation invariant, homogeneous of degree 0 function of $z_1, \dots, z_n$ (i.e. a function on the “residual moduli space”), then the weight:

$$h(x, z) = \tilde{\ell}_x \log f(z) = \ell_x \log f(z)$$

satisfies the functional equation (4.3) and is translation invariant and homogeneous of degree (-2) .

Prompted by these solutions, we want to get a global interpretation of the differential condition (4.3). Consider two simple paths $\gamma, \tilde{\gamma}$ in $\mathbb{H}$ started from x, y respectively, non intersecting and driven by smooth functions, for simplicity. The union of the two paths can be parametrized by Loewner’s equation in two extremal fashions: exploring γ first and then $\tilde{\gamma}$, or exploring $\tilde{\gamma}$ first and then γ . Define:

$$c(\gamma, z) = \int_0^\tau h(x_t, z_1(t), \dots, z_n(t)) dt$$

where γ has half-plane capacity 2τ , and $z_1(\cdot), \dots, z_n(\cdot)$ follow the Loewner flow driven by (x_t) . If $\gamma, \tilde{\gamma}$ have half-plane capacity $\varepsilon, c\varepsilon$ respectively, then:

$$c(\gamma, z) + c(\phi_\gamma(\tilde{\gamma}), \phi_\gamma(z)) = c(\tilde{\gamma}, z) + c(\phi_{\tilde{\gamma}}(\gamma), \phi_{\tilde{\gamma}}(z)) + O(\varepsilon^2).$$

This is exactly the content of (4.3), the corrections being as in the commutation relations. So it is easy to see (and straightforward to write, using comparisons of Loewner chains etc...) that for macroscopic paths $\gamma, \tilde{\gamma}$, we have:

$$c(\gamma, z) + c(\phi_\gamma(\tilde{\gamma}), \phi_\gamma(z)) = c(\tilde{\gamma}, z) + c(\phi_{\tilde{\gamma}}(\gamma), \phi_{\tilde{\gamma}}(z)).$$

Then we can define $c(\gamma \cup \tilde{\gamma}, z)$ to be this quantity. Similarly, one can grow a first half of γ , then a half of $\tilde{\gamma}$, then grow the end of $\gamma, \dots$, and get the same quantity. If A is any smooth hull (not intersecting the z_i 's), one can define $c(A, z) = c(\partial A, z)$, where ∂A is a smooth curve. Notice that this does not depend on the orientation of ∂A . Indeed, let x, y be the endpoints of ∂A , $\gamma, \tilde{\gamma}$ as above, such that γ and $\tilde{\gamma}$ are at distance at most ε of ∂A in the Hausdorff metric. Then $\phi_\gamma(\tilde{\gamma})$ and $\phi_{\tilde{\gamma}}(\gamma)$ have small half-plane capacity, and one can apply the previous result. If the boundary ∂A is described as the union of two arcs starting at x, y respectively, one gets the same result. Also, if A, B are two smooth hulls contained in a compact hull not intersecting the z_i 's, then, by similar arguments, we have the Lipschitz condition:

$$|c(A, z) - c(B, z)| \leq k(\text{cap}(A \cup B) - \text{cap}(A \cap B)).$$

Hence we can extend the definition of c by approximation by smooth hulls.

So if h is a function solving (4.3), then we can define a function $C = \exp c$ on hulls and residual configurations such that:

- (i). For all hulls A, B , $C(A.B, z) = C(B, \phi_A(z))C(A, z)$.
- (ii). If A is a hull of half-plane capacity ε located at x , then $C(A, z) = 1 + 2\varepsilon h(x, z) + o(\varepsilon)$.

Here $A.B$ designates the concatenation of the two hulls A, B : $\phi_{A.B} = \phi_B \circ \phi_A$. Conversely, if a nice function C satisfying (i) is given, we can recover a function h satisfying (4.3) (its derivative in the direction ℓ_x). So far, we have considered the following C 's:

$$C(A, z_1, \dots, z_{n+1}) = \prod_{i < j} \left(\phi'_A(z_i) \phi'_A(z_j) \left(\frac{z_j - z_i}{\phi_A(z_j) - \phi_A(z_i)} \right)^2 \right)^{\nu_{ij}} \frac{f(\phi_A(z_1), \dots, \phi_A(z_{n+1}))}{f(z_1, \dots, z_{n+1})}$$

where the marked point at infinity is now z_{n+1} . All the factors here are conformally invariant.

Let $\mathcal{H}$ designates a semigroup of hulls (as in [13]). The residual configuration is $(\mathbb{H}, z_1, \dots, z_{n+1} = \infty)$ (it no longer depends on the marked points x, y). More precisely, an element of $\mathcal{H}$ is a compact subset A of $\mathbb{H}$ such that $\mathbb{H} \setminus A$ is simply-connected and $\partial A \cap \mathbb{R} \subset \overline{A \cap \mathbb{H}}$; the semigroup is concatenation: $\phi_{A.B} = \phi_B \circ \phi_A$. Let $\mathcal{F}$ be a vector space of functions of the residual configuration $(z_1, \dots, z_{n+1})$ (say smooth functions). Then $\mathcal{H}$ acts on $\mathcal{F}$ by $A.f = f \circ \phi_A$ (note that this is not everywhere defined). If C is as above, C can be seen as a map $\mathcal{H} \rightarrow \mathcal{F}$ by $C(A)(z) = C(A, z)$. Then the condition (i) reads:

$$C(A.B) = C(A)(A.C(B))$$

which is saying that C is a 1-cocycle for (multiplicative) group cohomology for the $\mathcal{H}$ -module $\mathcal{F}$. This is formal since $\mathcal{H}$ is only a semigroup and the operation is not everywhere defined. Then the question is to determine the first cohomology group $H^1(\mathcal{H}, \mathcal{F})$, restricted to cocycles with Möbius covariance:

$$C(\phi(A), \phi(z)) = C(A, z)$$

where ϕ an homography. Let c_{ij} be the cocycle:

$$c_{ij}(A)(z) = \phi'_A(z_i) \phi'_A(z_j) \left(\frac{z_j - z_i}{\phi_A(z_j) - \phi_A(z_i)} \right)^2.$$

We have a natural map:

$$\begin{aligned} \mathbb{R}^{n(n+1)/2} &\longrightarrow H^1(\mathcal{H}, \mathcal{F}) \\ (\nu_{ij})_{i < j} &\longmapsto \prod_{i < j} (c_{ij})^{\nu_{ij}} \end{aligned}$$

The content of Section 8.5 in [13] is that this is an isomorphism when $n \leq 2$. At this point, it is not quite clear whether this is onto in general. Though it not one-to-one as soon as $n \geq 3$, since e.g. $c_{12}c_{34}/c_{13}c_{24}$ is a coboundary. It is easy to see that its image has dimension $n + 1$ and is generated for instance by $(c_{12}, c_{13}, \dots, c_{1,n+1}, c_{23})$.

So let us extend the discussion in [13] when $n \geq 3$. This involves several complications. The idea is to restrict to a subgroup of $\mathcal{H}$ (or rather a subalgebra of its tangent algebra) fixing a configuration $(z_1, \dots, z_{n+1})$. A cocycle restricts to a character of this subgroup, that must vanish on commutators. This gives differential conditions for tangent cocycles, that can then be integrated.

We begin by fixing three marked points at $0, 1, \infty$ (which we can do by a Möbius transformation). The other marked points are $z_1, \dots, z_{n-2}$. Consider the real Lie algebra generated by the Loewner fields:

$$A(x)\partial_w = \frac{w(w-1)}{w-x}\partial_w$$

that fix $0, 1, \infty$. Consider the subalgebra fixing the marked points $z_1, \dots, z_{n-2}$. This includes the fields:

$$A(x_1, \dots, x_{n-1}) = \det(A(x_i)(z_j))_{1 \leq i, j \leq n-1} = \prod_{i=1}^{n-1} z_i(z_i - 1) \frac{V(x_1, \dots, x_{n-1})V(z_1, \dots, z_{n-1})}{\prod_{i,j} z_i - x_j}$$

with the convention $w = z_{n-1}$, where V is the Vandermonde polynomial $V(x_1, \dots, x_{n-1}) = \prod_{i < j} (x_j - x_i)$ (Cauchy determinant). By taking limits of such vector fields (as $x_1, \dots, x_n \rightarrow x$, normalizing by $V(x_1, \dots, x_n)$), one gets $\hat{A}(x) = (n-2)!P(w)/(w-x)^{n-1}$, where $P(w) = w(w-1)(w-z_1) \cdots (w-z_{n-2})$ is a polynomial of degree n . The (tangent) cocycle dC restricted to this subalgebra is a morphism to the (trivial) Lie algebra $\mathbb{R}$, so it vanishes on commutators. By considering the limit of $[\hat{A}(x), \hat{A}(y)]$ as $y \searrow x$, we see that $\hat{A}(x) = (2n-1)!P(w)^2/(w-x)^{2n}$ is annihilated by dC . It is easy to see that:

$$\hat{A}(x) = (\partial_x^{2n-1} P(x) \partial_x^{2-n}) \tilde{A}(x).$$

Note that $(\partial_x^{2n-1} P(x) \partial_x^{2-n})$ is a well-defined linear differential operator of degree $n+1$, with polynomial coefficients. It follows that:

$$dC(\tilde{A}(x)) = a + \frac{a_0}{x^{n-1}} + \frac{a_1}{(x-1)^{n-1}} + \sum_{j=1}^{n-2} \frac{b_j}{(x-z_j)^{n-1}}$$

where the coefficients depend on the marked points $z_1, \dots, z_{n-2}$. We want to integrate this to get information on $dC(A(x_1, \dots))$. Consider the atomic measure (in the variable x):

$$m = \sum_i (-1)^{n-1-i} V(x_1, \dots, \hat{x}_i, \dots) \delta_{x_i}.$$

One can think of m as the Vandermonde determinant $V(x_1, \dots, x_{n-1})$ where the last row $(x_1^{n-2}, \dots, x_{n-1}^{n-2})$ is replaced by $(\delta_{x_1}, \dots, \delta_{x_{n-1}})$. Then we have the decomposition:

$$\frac{V(x_1, \dots, x_{n-1})}{\prod_i (w - x_i)} = \left\langle m, \frac{1}{w-x} \right\rangle = (-1)^n \left\langle m^{(2-n)}, \frac{(n-2)!}{(w-x)^{n-1}} \right\rangle$$

where $m^{(2-n)}$ can be chosen with a continuous, compactly supported density (since $\langle m, 1 \rangle = \dots = \langle m, x^{n-3} \rangle = 0$, which follows from the Vandermonde form of m). It follows that:

$$A(x_1, \dots, x_{n-1}) = (-1)^n \prod_{i=1}^{n-2} z_i(z_i - 1) \frac{V(z_1, \dots, z_{n-2})}{\prod_{i < n-1, j \leq n-1} z_i - x_j} \langle m^{(2-n)}, \tilde{A}(x) \rangle$$

and consequently:

$$dC(A(x_1, \dots, x_{n-1})) = \prod_{i=1}^{n-2} z_i(z_i - 1) \frac{V(z_1, \dots, z_{n-2})}{\prod_{i < n-1, j \leq n-1} z_i - x_j} \left\langle m, ax^{n-2} + \frac{a_0}{x} + \frac{a_1}{x-1} + \sum_{j=1}^{n-2} \frac{b_j}{x - z_j} \right\rangle$$

for some coefficients a, a_0, b_j depending on $z_1, \dots, z_{n-2}$. From the Vandermonde form of m and simple manipulations, we get:

$$dC(A(x_1, \dots, x_{n-1})) = \prod_{i=1}^{n-2} z_i(z_i - 1) \frac{V(z_1, \dots, z_{n-2})V(x_1, \dots, x_{n-1})}{\prod_{i < n-1, j \leq n-1} z_i - x_j} \left(a + \frac{a_0}{\prod_j x_j} + \frac{a_1}{\prod_j x_j - 1} + \sum_{k=1}^{n-2} \frac{b_k}{\prod_j x_j - z_k} \right).$$

Using the Cauchy determinant formula (and its derivative w.r.t z_k), it is then easy to see that:

$$\begin{vmatrix} \frac{1}{x_1 - z_1} & \dots & \frac{1}{x_{n-1} - z_1} \\ \vdots & & \vdots \\ \frac{1}{x_1 - z_{n-2}} & \dots & \frac{1}{x_{n-1} - z_{n-2}} \\ R(x_1)(z) & \dots & R(x_{n-1})(z) \end{vmatrix} = 0$$

where:

$$R(x) = dC(A(x)) - \left(a + \frac{a_0}{x} + \frac{a_1}{x-1} + \sum_{j=1}^{n-2} \frac{b_j}{(x - z_j)^2} \right)$$

which depends implicitly on the z variables (and we have replaced a_0 with $a_0 / \prod_i z_i$, a_1 with $a_1 / \prod_i (z_i - 1)$; since we use 0,1 for normalization, this induces some asymmetry). For the part corresponding to the coefficient a , one can consider the limit $w \rightarrow \infty$ of $A(x_1, \dots, x_{n-1})$. If $Q(x) = (x - z_1) \dots (x - z_{n-2})$, after row operations, we get:

$$\begin{vmatrix} 1 & \dots & 1 \\ \vdots & & \vdots \\ x_1^{n-3} & \dots & x_{n-1}^{n-3} \\ Q(x_1)R(x_1)(z) & \dots & Q(x_{n-1})R(x_{n-1})(z) \end{vmatrix} = 0$$

If we apply $\partial_{x_1}^{n-2}$ to this expression (using multilinearity in rows), we get $\partial_{x_1}^{n-2} Q(x_1)R(x_1)(z) = 0$. Hence:

$$dC(A(x)) = a + \frac{a_0}{x} + \frac{a_1}{x-1} + \sum_{j=1}^{n-2} \frac{c_j}{x - z_j} + \sum_{j=1}^{n-2} \frac{b_j}{(x - z_j)^2}$$

where coefficients may depend on z .

In terms of h , this means that:

$$h(x, z) = \sum_i \frac{\mu_i(z)}{(x - z_i)^2} + \sum_i \frac{\rho_i(z)}{x - z_i}$$

where μ_i, ρ_i are translation invariant and homogeneous of degree 0, -1 respectively (for normalization reasons, we have to divide $dC(A(x))$ by $x(x-1)$). So we can substitute this expression in the commutation equation (4.3), and get after some simplifications:

$$\sum_{i,j} \left(\frac{1}{(y - z_j)(x - z_i)^2} - \frac{1}{(x - z_j)(y - z_i)^2} \right) \partial_j \mu_i + \left(\frac{1}{(y - z_j)(x - z_i)} - \frac{1}{(x - z_j)(y - z_i)} \right) \partial_j \rho_i = 0$$

as a rational function in x, y . This implies that $\partial_j \mu_i = 0$ for all i, j (considering first the coefficient of $(x - z_i)^{-2}$, and then letting y vary). Similarly, considering the coefficient of $(x - z_i)^{-1}$, we get:

$$\sum_j \frac{\partial_j \rho_i - \partial_i \rho_j}{y - z_j} = 0$$

and letting y vary, we get the cross-derivative condition $\partial_i \rho_j = \partial_j \rho_i$. Hence we can find constant coefficients μ_i and a function $f(z)$ such that:

$$h(x, z) = \sum_i \frac{\mu_i}{(x - z_i)^2} + \ell_x \log f(z).$$

Note that $f(z) = (z_j - z_i)^{\nu_{ij}}$ produces the term $2\nu_{ij}/(x - z_i)(x - z_j)$. So we can rewrite h as:

$$h(x, z) = \sum_{i,j} \nu_{ij} \left(\frac{1}{x - z_i} - \frac{1}{x - z_j} \right)^2 + \ell_x \log f(z)$$

in a non-unique fashion (e.g. take f the power of a cross-ratio; here $z_{n+1} = \infty$). Since:

$$\partial_j \sum_i \rho_i = \left(\sum_i \partial_i \right) \rho_j = 0, \quad \partial_j \sum_i z_i \rho_i = \left(\sum_i z_i \partial_i \right) \rho_j + \rho_j = 0$$

f is translation invariant and homogeneous of some fixed degree. This degree can be set to 0 by adjusting the ν_{ij} 's. One can freely set all the ν_{ij} 's to zero except $(\nu_{12}, \dots, \nu_{1,n+1}, \nu_{23})$ e.g., which gives $(n + 3)$ numerical invariants.

Let us sum up the previous discussion.

Lemma 4. *If the system*

$$\begin{cases} \frac{\kappa}{2} \partial_{xx} \psi + \sum_i \frac{2\partial_i \psi}{z_i - x} + \frac{2\partial_y \psi}{y - x} + \left(\left(1 - \frac{6}{\tilde{\kappa}} \right) \frac{1}{(y - x)^2} + h_1(x, z) \right) \psi = 0 \\ \frac{\tilde{\kappa}}{2} \partial_{yy} \psi + \sum_i \frac{2\partial_i \psi}{z_i - y} + \frac{2\partial_x \psi}{x - y} + \left(\left(1 - \frac{6}{\kappa} \right) \frac{1}{(x - y)^2} + h_2(y, z) \right) \psi = 0 \end{cases}$$

admits a non-vanishing solution ψ (smooth, homogeneous and translation invariant), then:

(i). *Either $\tilde{\kappa} = \kappa$ or $\tilde{\kappa} = 16/\kappa$.*

(ii). *The functions h_1, h_2 can be written as $h_1(x, z) = h(x, z)$, $h_2(y, z) = h(y, z)$, where:*

$$h(x, z) = \sum_{1 \leq i < j \leq n+1} 2\nu_{ij} \left(\frac{1}{x - z_i} - \frac{1}{x - z_j} \right)^2 + \ell_x \log f(z)$$

where ν_{ij} are constant parameters and $f(z)$ is a conformally invariant function of the marked points $(z_1, \dots, z_{n+1})$.

Moreover the weight vector $(\nu_{ij})_{i < j}$ is well-defined modulo the relations $\nu_{ij} + \nu_{kl} = \nu_{il} + \nu_{jk}$.

Conversely, it is natural to ask whether these conditions are sufficient for the existence of a non vanishing solution ψ . In the situation where h is the rational function:

$$h(x, z) = \sum_i \frac{\mu_i}{(x - z_i)^2} + \sum_{i < j} \frac{\nu_{ij}}{(x - z_i)(x - z_j)}$$

then we can find an elementary solution of the form:

$$\psi(x, y, z) = \left(\prod_i (z_i - x)^{a_i} (z_i - y)^{\tilde{a}_i} \right) (y - x)^b \prod_{i < j} (z_j - z_i)^{b_{ij}}$$

If $\tilde{\kappa} = \kappa$, we can pick $b = 2/\kappa$, $\tilde{a}_i = a_i$, $b_{ij} = \nu_{ij}/2$ and a_i a solution of $(\kappa/2)a_i(a_i - 1) + 2a_i + \mu_i = 0$ for all i . If $\tilde{\kappa} = 16/\kappa$, set $b = -1/2$, $\tilde{a}_i = -a_i\kappa/4$, $b_{i,j} = \nu_{ij}/2$ and a_i a solution of $(\kappa/2)a_i(a_i - 1) + 2a_i + \mu_i = 0$ for all i . Finding ψ of a prescribed homogeneity degree is much more difficult.

For a general h , we shall later discuss martingale interpretations of solutions. If one considers n SLEs, $n \geq 2$, one gets a system of n linear PDEs (with coefficients $h_1, \dots, h_n$ to be specified). We give an example of this situation in the next section.

4.2 Restriction, locality and cocycles

In the previous subsection, we made some progress on the question of classifying restriction measures, which we now explicit (in the chordal setting). A configuration is a simply connected domain D with n marked points on the boundary: $(D, z_1, \dots, z_n)$ (say D has Jordan boundary). The following definition is a natural extension of [13]. We call a restriction measure a collection μ of measures parametrized by the configuration $(D, z_1, \dots, z_n)$ such that:

- (i). The measure $\mu_{(D, z_1, \dots)}$ is supported on simply- connected compact subsets of the compactification of D that intersect the boundary of D exactly at $z_1, \dots, z_n$. Also, $\partial K \cap \partial D \subset \overline{K \cap D}$.
- (ii). (Conformal invariance) If $\phi : (D, z_1, \dots) \rightarrow (D', z'_1, \dots)$ is an equivalence of configurations, $\phi_* \mu_{(D, \dots)} = \mu_{(D', \dots)}$.
- (iii). (Restriction property) If $(D, z_1, \dots)$ and $(D', z_1, \dots)$ are configurations such that $D' \subset D$ and $\partial D' \cap \partial D$ contains neighbourhoods of $z_1, \dots, z_n$, then: $\mu_D(\cdot| \subset D') = \mu_{D'}$.

Given such a restriction measure (strictly speaking, collection of measures, or measure-valued function on the moduli space), we can define a cocycle: $C_\mu(A, z) = \mu_{(\mathbb{H}, z)}(\{K : K \cap A = \emptyset\})$. Indeed, the tower property of conditional expectations:

$$\mu_{(\mathbb{H}, z)}(\cdot| \subset \mathbb{H} \setminus A.B) = \mu_{(\mathbb{H}, z)}(\cdot| \subset \mathbb{H} \setminus A.B| \subset \mathbb{H} \setminus A) = \mu_{(\mathbb{H} \setminus A, z)}(\cdot| \subset \mathbb{H} \setminus A.B) = \mu_{(\mathbb{H}, A, z)}(\cdot| \subset \mathbb{H} \setminus B)$$

translates into the cocycle condition: $C_\mu(A.B, z) = C_\mu(A, z)C_\mu(B, A.z)$. Besides, μ is entirely determined by C_μ , since the events $\{K \cap A = \emptyset\}$ constitute a π -system generating the full σ -algebra for random compact sets K satisfying the topological condition (i).

We have proved that under a regularity assumption C_μ can be expressed as:

$$C_\mu(A, z) = \prod_{i < j} \left(\phi'_A(z_i) \phi'_A(z_j) \left(\frac{z_j - z_i}{\phi_A(z_j) - \phi_A(z_i)} \right)^2 \right)^{\nu_{ij}} \frac{f(\phi_A(z_1), \dots, \phi_A(z_n))}{f(z_1, \dots, z_n)}.$$

The question is what cocycles can be realized through a restriction measure. A result of [13] is that if $n = 2$, then the cocycle $\left(\phi'_A(z_1) \phi'_A(z_2) \left(\frac{z_2 - z_1}{\phi_A(z_2) - \phi_A(z_1)} \right)^2 \right)^\nu$ corresponds to a restriction measure iff $\nu \geq 5/8$. Also, there is an operation $\bullet$ on restriction measures (filling of the union of independent samples) such that $C_{\mu_1 \bullet \mu_2} = C_{\mu_1} C_{\mu_2}$.

Let us now discuss locality; the close relation between locality and restriction is stressed in [13]. Consider a configuration (D, a, b, c) with three marked points on the boundary, in cyclic order. Using chordal SLE₆, one can define a distribution μ relative to such configurations and satisfying:

- (i). $\mu_{(D,a,b,c)}$ is supported on simply connected compact subsets K of the compactification of D whose intersection with ∂D consists in a point $X \in (bc)$ and an arc contained in (cb) and containing a . Also, $\partial K \cap \partial D \subset \overline{K \cap D}$.
- (ii). (Conformal invariance) If $\phi : (D, a, b, c) \rightarrow (D', a', b', c')$ is an equivalence of configurations, then $\phi_* \mu_{(D,a,b,c)} = \mu_{(D',a',b',c')}$.
- (iii). (Locality) Let $\mu_{(D,a,b,c,x)}$ be the disintegrated measure $\mu_{(D,a,b,c)}(\cdot | X \in dx)$. If (D, a, b, c) and (D', a, b', c') are configurations such that $D' \subset D$, $b' \in (ab)$, $c' \in (ca)$, $\partial D' \cap \partial D$ contains the arcs (ab') , $(c'a)$ and a neighbourhood of x , then:

$$\mu_{(D,a,b,c,x)}(\cdot | K \subset D', \partial K \subset (ab') \cup \{x\} \cup (c'a)) = \mu_{(D',a,b',c',x)}.$$

Let us associate a cocycle to this collection of measures. We phrased the locality property as a restriction property for disintegrated measures, so we can define as above:

$$C(A, z) = \mu_{(\mathbb{H}, a, b, c, x)}(\{K : K \subset \mathbb{H} \setminus A\})$$

where $z = (a, b, c, x)$. With these conditions, the measures are entirely determined by the distribution of X . More precisely, if this distribution is $f(x)dx$ for the configuration (D, a, b, c) and ϕ is the equivalence $(D, a, b, c) \rightarrow (D', a, b', c')$, then $C_\mu(A, z) = \phi'(x)f \circ \phi(x)/f(x)$, or, without assumptions on the normalization of the equivalence ϕ :

$$C_\mu(\phi, z) = \frac{c_{ax}c_{bc}^{1/2}}{c_{ab}^{1/2}c_{ac}^{1/2}}(\phi, z) \frac{f \circ \phi(z)}{f(z)}$$

and again one can recover μ from C_μ . One can think of several extensions to configurations with more points. Consider the following example, coming from percolation. For critical percolation in a rectangle $(ABCD)$, consider the highest (resp. lowest) open path from (AB) to (CD) with endpoints A', B', C', D' respectively. This gives a measure (with mass < 1) on hulls delimited by the two paths. Disintegrated measures (w.r.t. $A', \dots, D'$) have the restriction property. The measure on hulls is determined by the joint distribution (A', B', C', D') , than can be obtained by taking partial derivatives of Cardy's formula (see [9] for related questions).

5 A particular case

In this section, we discuss the important situation where all marked points on the boundary are growth points for commuting SLEs. This situation is studied in greater details in [10].

As described earlier, consider the half-plane $\mathbb{H}$ with $(2n)$ marked points on the boundary, $(x_1, \dots, x_{2n})$, in cyclical order. Consider n independent $\text{SLE}_{8/3}$ from x_{2i-1} to x_{2i} , $i = 1, \dots, n$. Define $\psi(x_1, \dots, x_{2n})$ to be the probability of no pairwise intersection. This function is invariant under the full Möbius group (∞ is only used for normalization). Now, if γ_i is the trace of the i -th SLE, define

$$\begin{aligned} \psi(x_i) &= \mathbb{P}(\gamma_i \cap \gamma_j = \emptyset, 1 \leq i < j \leq n) \\ &= \mathbb{P}(\gamma_1(0, t) \cap (\cup_{j>1} \gamma_j) = \emptyset, \gamma_1(t, \infty) \cap (\cup_{j>1} \gamma_j) = \emptyset, \gamma_i \cap \gamma_j = \emptyset, 2 \leq i < j \leq n) \\ &= \prod_{j>1} \mathbb{P}(\gamma_j \cap \gamma_1(0, t) = \emptyset) \mathbb{P}(\tilde{\gamma}_i \cap \tilde{\gamma}_j = \emptyset, 1 \leq i < j \leq n) \end{aligned}$$

where $\tilde{\gamma}_i$ are independent $\text{SLE}_{8/3}$ in the domain $(\mathbb{H}, g_t(\gamma_1(t)), g_t(x_2), \dots, g_t(x_{2n}))$ and (g_t) are the conformal equivalences associated with the first SLE. This relies on the Markov property for γ_1 , the restriction property for each γ_j , $j > 1$, and induction on n . As a consequence, the following process:

$$\psi(W_t, g_t(x_2), \dots, g_t(x_{2n})) \prod_{j>1} \left(g'_t(x_{2j-1}) g'_t(x_{2j}) \left(\frac{x_{2j} - x_{2j-1}}{g_t(x_{2j}) - g_t(x_{2j-1})} \right)^2 \right)^{\alpha_\kappa}$$

is a martingale, where $\alpha_\kappa = (6 - \kappa)/2\kappa$, $\kappa = 8/3$. Now, one can do this starting at each point x_i (since $\text{SLE}_{8/3}$ is reversible). This implies that ψ is annihilated by the operators ($k = 1, \dots, 2n$):

$$\frac{\kappa}{2}\partial_{kk} + \sum_{l \neq k} \frac{2\partial_l}{x_l - x_k} + (\kappa - 6) \frac{\partial_k}{x_k - x_{\iota(k)}} + \frac{\kappa - 6}{\kappa} \sum_{\{j, \iota(j)\} \neq \{k, \iota(k)\}} \left(\frac{1}{x_j - x_k} - \frac{1}{x_{\iota(j)} - x_k} \right)^2$$

where ι defines the chosen pairing $\iota(2i - 1) = 2i$, $\iota(2i) = 2i - 1$.

This is not very symmetrical. It is easy to see that the function

$$\psi(\dots x_i \dots) \prod_j (x_{2j} - x_{2j-1})^{1-6/\kappa}$$

is annihilated by the operators:

$$\begin{cases} \frac{\kappa}{2}\partial_{k,k} + \sum_{l \neq k} \frac{2\partial_l}{x_l - x_k} + \frac{\kappa - 6}{\kappa} \sum_{l \neq k} \frac{1}{(x_l - x_k)^2}, & k = 1, \dots, 2n \\ \sum_k \partial_k \\ \sum_k x_k \partial_k - n(1 - 6/\kappa) \\ \sum_k x_k^2 \partial_k - (1 - 6/\kappa)(x_1 + \dots + x_{2n}) \end{cases}$$

the last three ones corresponding to the invariance of ψ under the Moebius group. In fact, as is discussed in [10], one can make sense of this system for any $\kappa \in (0, 8/3)$ using appropriate loop-soups.

Let us make a few remarks on this system. First, each choice of a non-crossing pairing of the $(2n)$ boundary points yields a solution; there are C_n such pairings. If $\kappa = 6$, this is the system satisfied by crossing probabilities for critical percolation in a $(2n)$ -gon with alternating boundary conditions. The number of these crossing probabilities is the number of non-crossing partitions of the set of blue edges, which is known to be C_n . In the case $n = 2$, it is trivial to solve this system, which reduces to a hypergeometric equation (and $C_2 = 2$). If $n = 3$, one can write this system in a Pfaffian form, proving that its rank is indeed $C_3 = 5$. In the case $\kappa = 6$, $n = 3$, and configurations with 3-fold symmetry, one can express solutions in terms of ${}_3F_2$. Finally, one can take the limit $\kappa \rightarrow \infty$ of the system; in this case, solutions are polynomials, and it is easy to see that the rank of the system is C_n for all n . Euler integrals for solutions of this system are discussed in [10].

6 Local commutation

In this section we see how to go from infinitesimal commutation relations to commutation (in law) of SLE hulls. Recall from Section 3 the definition of an $\text{SLE}(\kappa, b, \tilde{\kappa}, \tilde{b})$. We have seen in the previous sections that the existence of such an SLE implies conditions on $(\kappa, b, \tilde{\kappa}, \tilde{b})$ (in particular either $\tilde{\kappa} = \kappa$ or $\tilde{\kappa} = 16/\kappa$). Conversely, assume that the data $(\kappa, b, \tilde{\kappa}, \tilde{b})$ satisfies the appropriate conditions. We will see that this implies the existence of an $\text{SLE}(\kappa, b, \tilde{\kappa}, \tilde{b})$. Note that this is not saying anything on the long time behaviour of such an SLE. The questions involving collisions of commuting SLEs are delicate and cannot be handled by these methods.

Proposition 5. *Consider the upper half-plane $\mathbb{H}$ with $n + 2$ distinct marked real points $x, y, z_1, \dots, z_n$. Assume that $\kappa, \tilde{\kappa}$ are positive numbers and $b, \tilde{b}$ are smooth functions, translation invariant and homogeneous of degree (-1) , such that the following relation holds:*

$$[\mathcal{L}, \mathcal{M}] = \frac{4}{(y - x)^2} (\mathcal{M} - \mathcal{L})$$

where $\mathcal{L}$ (resp. $\mathcal{M}$) is the infinitesimal generator of $\text{SLE}_\kappa(b)$ (resp. $\text{SLE}_{\tilde{\kappa}}(\tilde{b})$) growing at x (resp. y). Then there exists an $\text{SLE}(\kappa, b, \tilde{\kappa}, \tilde{b})$.

We will use the following lemma. Let ϕ_0 be some conformal equivalence $\mathbb{H} \setminus K \rightarrow \mathbb{H}$, with hydrodynamic normalization at infinity. Let us call ϕ_0 -capacity the increasing function on hulls: $\text{cap} \circ \phi_0^{-1} - \text{cap}(K)$.

Lemma 6. *With the hypotheses of the Proposition, let D_1, D_2 be disjoint compact neighbourhoods of x, y resp., with Jordan boundary, not containing any other marked point; η_1, η_2 are two positive numbers. Then the two following procedures define the same probability law on pairs of chains in $\mathbb{H}$:*

- (i). *Grow an $\text{SLE}_\kappa(b)$ at x until it exits D_1 or its ϕ_0 -capacity exceeds η_1 , Then grow an independent $\text{SLE}_{\tilde{\kappa}}(\tilde{b})$ at y in the remaining domain, until it exits D_2 or its ϕ_0 -capacity exceeds η_2 .*
- (ii). *Grow an $\text{SLE}_{\tilde{\kappa}}(\tilde{b})$ at y until it exits D_2 or its ϕ_0 -capacity exceeds η_2 . Then grow an independent $\text{SLE}_\kappa(b)$ at x in the remaining domain, until it exits D_1 or its ϕ_0 -capacity exceeds η_1 .*

Proof. Informally, the argument is the following: divide the two SLEs in n segments; one has to prove that one can either grow the n segments of the first SLE, then the n segments of the second SLE, or the other way round and get the same law. The permutation of two segments (of the two SLEs) induces an error term of $O(n^{-3})$, from the infinitesimal commutation relations. One needs n^2 such permutations; letting n go to infinity, one gets the result. The uniformity in the error terms is provided by the restriction to paths in the disjoint compact neighbourhoods D_1, D_2 .

For simplicity, we will consider only the case where $\phi_0 = \text{Id}$ (and the ϕ_0 -capacity is the ordinary half-plane capacity). For the general case, one has to replace fixed times by corresponding stopping times; the proof goes otherwise unchanged.

For positive times S, T , let $\mathbb{E}_1$ designate the expectation for pairs of random curves obtained by growing first the SLE in D_1 up to time S (half-plane capacity $2S$), and then the SLE in D_2 up to time T . The symbol $\mathbb{E}_2$ refers to expectation for the reversed construction. The driving process for each of these Loewner chains (seen in the original half-plane) is denoted by X, Y . Let τ_1 be the time at which the first SLE exits D_1 , and τ_2 the corresponding time for the second SLE. We will prove that

$$\mathbb{E}_1(\mathbf{1}_{\tau_1 > S, \tau_2 > T}) = \mathbb{E}_2(\mathbf{1}_{\tau_1 > S, \tau_2 > T})$$

as measures on $C_0([0, S], \mathbb{R}) \times C_0([0, T], \mathbb{R})$. To recover the statement of the lemma, one then considers the measures:

$$\mathbb{E}_i(\mathbf{1}_{\tau_1 > S, \tau_2 > T} - \mathbf{1}_{\tau_1 > S+ds, \tau_2 > T})$$

for $i = 1, 2$. So we can work with fixed times S and T . Note that $2\tau_i$ is bounded by the half-plane capacity of D_i , $i = 1, 2$.

Let $S_0 = 0 < S_1 < \dots < S_m = S$ and $T_0 = 0 < T_1 < \dots < T_m = T$ be fixed sequences of times ($m \geq 1$). Also, let $(\tilde{\varphi}_i)_{0 \leq i \leq m}, (\tilde{\psi}_i)_{0 \leq i \leq m}$ be test functions (i.e. in $C_c^\infty(\mathbb{R})$). By a monotone class argument, we need only to see that:

$$\mathbb{E}_1 \left(\prod_k \tilde{\varphi}_k(X_{S_k}) \tilde{\psi}_k(Y_{T_k}) \mathbf{1}_{\tau_1 > S, \tau_2 > T} \right) = \mathbb{E}_2 \left(\prod_k \tilde{\varphi}_k(X_{S_k}) \tilde{\psi}_k(Y_{T_k}) \mathbf{1}_{\tau_1 > S, \tau_2 > T} \right)$$

For $n \geq 1$, consider increasing sequences $(s_i)_{0 \leq i \leq mn}, (t_i)_{0 \leq i \leq mn}$, where $s_{nj} = S_j, t_{nj} = T_j$, and the increments $(s_{i+1} - s_i), (t_{i+1} - t_i)$ go uniformly to 0. Define $\varphi_{ni} = \tilde{\varphi}_i, \psi_{ni} = \tilde{\psi}_i$, and $\varphi_i = \psi_i = 1$ if n does not divide i .

Note that the commutation relation holds for functions of the positions of all marked points in the Loewner flow. For convenience, we will approximate the event $\{\tau_1 > S, \tau_2 > T\}$ by a function of an extended flow. More precisely, let $\delta > 0$ be a (small) positive number and N a (large) integer. Mark N points $z_1, \dots, z_N$ on the Jordan boundaries of D_1 and D_2 (one can also mark their conjugates $\bar{z}_1, \dots, \bar{z}_N$, extending the flow

by Schwarz reflection). For instance, one can choose them so that the Hausdorff distance between $\partial D_1 \cup \partial D_2$ and $\{z_1, \dots, z_N\}$ is minimal (N being fixed).

Let $\mathcal{K} = \{(K, u)\}$ where K is a compact hull included in D_1 and u is a point in $\partial D_1 \cap K$. Then $\mathcal{K}$ is a compact set (using the Hausdorff metrics on compact subsets of D_1), so one can choose $\delta > 0$ so that:

$$\delta > \max_{(K, u) \in \mathcal{K}} \left(\min_i |\phi_K(z_i) - \phi_K(u)| \right)$$

and the corresponding inequality holds for hulls in D_2 . Also, it is easy to see that one can choose δ so that it goes to zero as N goes to infinity, by a compactness argument. Let f_δ be a smooth function of the variables $(x, y, z_1, \dots, z_N)$, taking values in $[0, 1]$, such that it vanishes if $|x - z_i| < \delta$ or $|y - z_i| < \delta$ for some $i \in \{1, \dots, N\}$ and equals 1 if $|x - z_i| > 2\delta$, $|y - z_i| > 2\delta$ for all i . One can assume that one of the z_i 's is real and between x and y , and similarly, the other marked points (that influence the drift) are separated from x , y by one of the "spectator" z_i 's. The choice of δ ensures that f_δ vanishes as soon as an SLE crosses ∂D_i , $i = 1, 2$.

Let W denote the full configuration (images of growth points and marked points in the Loewner flow). Then we just have to prove that:

$$\lim_{n \rightarrow \infty} (\mathbb{E}_1 - \mathbb{E}_2) \left(\prod_k \varphi_k(X_{s_k}) f_\delta(W_{(s_k, 0)}) \psi_k(Y_{t_k}) f_\delta(W_{(0, t_k)}) \right) = 0$$

and then let $N \nearrow \infty$, $\delta \searrow 0$ to get the result for stopped SLEs. So it what follows we may replace $\varphi_k(X_{s_k}) f_\delta(W_{(s_k, 0)})$ with $\varphi_k(W_{(s_k, 0)})$ (a function of the configuration), and similarly $\psi_k(X_{s_k}) f_\delta(W_{(s_k, 0)})$ with $\psi_k(W_{(s_k, 0)})$.

Consider also two random curves $\gamma, \hat{\gamma}$ started from x (resp. y) in $\mathbb{H}$, parameterized by half-plane capacity. Let $\phi_{s,t} = \phi_{\gamma_{[0,s]} \cup \hat{\gamma}_{[0,t]}}$, and $W_{s,t}$ is the configuration $\phi_{s,t}(\gamma_s, \hat{\gamma}_t, z_1, \dots, z_N, \dots)$. We will also abbreviate $x_k = W_{(s_k, 0)}$, $y_k = W_{(0, t_k)}$.

Consider two permutations σ and σ' of $\{s_1, \dots, s_{mn}, t_1, \dots, t_{mn}\}$, increasing for the partial order generated by $s_k < s_{k+1}$, $t_k < t_{k+1}$, and such that σ and σ' differ by a transposition of two consecutive elements. For instance $\sigma = (s_1, \dots, s_{mn}, t_1, \dots, t_{mn})$ and $\sigma' = (s_1, \dots, s_{mn-1}, t_1, s_{mn}, t_2, \dots, t_{mn})$. Suppose that $\gamma, \hat{\gamma}$ are obtained from the permutation σ in the following fashion: if $\sigma = (\sigma_1, s_{i+1}, \sigma_2)$, t_j is the maximal t . element in σ_1 , and $\phi = \phi_{s_i, t_j}$, then $\phi(\gamma_{[s_i, s_{i+1}]})$ is an $\text{SLE}_\kappa(b)$ started from W_{s_i, t_j} and independent of ϕ conditionally on its starting state (stopped so that $\phi_{s_i, 0}(\gamma_{[s_i, s_{i+1}]})$ has capacity $2(s_{i+1} - s_i)$). Likewise, if $\sigma = (\sigma_1, t_{j+1}, \sigma_2)$, s_i is the maximal s . element in σ_1 , and $\phi = \phi_{s_i, t_j}$, then $\phi(\gamma_{[t_j, t_{j+1}]})$ is an $\text{SLE}_{\tilde{\kappa}}(\tilde{b})$ started from W_{s_i, t_j} and independent of ϕ conditionally on its starting state. The symbol $\mathbb{E}$ is expectation for this construction (relative to σ), and $\mathbb{E}'$ is the corresponding expectation obtained from σ' .

Let $\sigma = (\sigma_1, s_i, t_j, \sigma_2)$ and $\sigma' = (\sigma_1, t_j, s_i, \sigma_2)$. Then:

$$\mathbb{E}' \left(\prod_k \varphi_k(x_k) \psi_k(y_k) \right) = \mathbb{E}' \left(\left(\prod_{k < i, l < j} \varphi_k(x_k) \psi_l(y_l) \right) \varphi_i(x_i) \psi_j(y_j) \left(\prod_{k > i, l > j} \varphi_k(x_k) \psi_l(y_l) \right) \right)$$

(Here the f_δ are implicitly included in the φ_k, ψ_k). The expectation of the last part of the product conditionally on $\gamma_{[0, s_i]}, \hat{\gamma}_{[0, t_j]}$ is a function of W_{s_i, t_j} . Denote by $F(u, v)$ this function, which is the same under $\mathbb{E}$ and $\mathbb{E}'$; by induction and standard regularity results (the drift terms stay bounded as long as the functional does not vanish), it is easily seen that F is a smooth function; the existence of regular conditional probability is clear for the same reasons.

Now, consider:

$$\mathbb{E} \left(\varphi_i(x_i) \psi_j(y_j) F(\phi_{s_i, t_j}(\gamma_{s_i}, \hat{\gamma}_{t_j})) \middle| W_{s_{i-1}, t_{j-1}} \right) - \mathbb{E}' \left(\varphi_i(x_i) \psi_j(y_j) F(\phi_{s_i, t_j}(\gamma'_{s_i}, \hat{\gamma}'_{t_j})) \middle| W_{s_{i-1}, t_{j-1}} \right)$$

Assume that i, j are not multiples of n (and traces are away from the boundaries of D_1, D_2 at time i, j). Then $\varphi_i = \psi_j = 1$, and this difference is $O(n^{-3})$, from the infinitesimal commutation relation.

If i or j is a multiple of n , note that, if $x' = \phi_{s_i, t_{j-1}}(\gamma_{s_i})$, $y' = \phi_{s_{i-1}, t_j}(\gamma_{t_j})$, $x'' = \phi_{s_i, t_j}(\gamma_{s_i})$, $y'' = \phi_{s_i, t_j}(\gamma_{t_j})$, then:

$$x' = x'' - \frac{2(t_j - t_{j-1})}{x'' - y''} + O(n^{-2})$$

using the backward Loewner flow; one gets a similar expression for y' , and these hold under $\mathbb{E}$ and $\mathbb{E}'$. So in this case the difference is $O(n^{-2})$.

To get from $\sigma = (s_1, \dots, s_{mn}, s_1, \dots, s_{mn})$ to $\sigma' = (t_1, \dots, t_{mn}, t_1, \dots, t_{mn})$, one needs $(mn)^2$ transpositions ((mn) transpositions to bring t_1 in first position, then (mn) transpositions to bring t_2 in second position, ...). For such a transposition (s_i, t_j) , i or j is a multiple of n in $m^2(2n-1)$ case. This transposition is valid as long as the paths stay in D_1, D_2 (more precisely, as long as the f_δ terms are 1). Conversely, if a path is close to the boundary of D_1, D_2 , the functional is zero with probability close to one. Hence:

$$(\mathbb{E}_1 - \mathbb{E}_2) \left(\prod_k \varphi_k(X_{s_k}) \psi_k(Y_{t_k}) \mathbf{1}_{E(n, N, \delta)} \right) = m^2(n-1)^2 O(n^{-3}) + m^2(2n-1) O(n^{-2}) = O(C(N, \delta)/n)$$

where $X_s = \phi_{s,0}(\gamma_s)$, $Y_t = \phi_{0,t}(\hat{\gamma}_t)$, and $E(n, N, \delta)$ is the event that none of the f_δ 's vanishes at a sampled time. The error term is uniform in n but depends on N, δ .

As n goes to infinity (N, δ being fixed), the probability that the first SLE crosses ∂D_1 without f_δ vanishing at one of the sampled times s_i goes to zero (since in this case f_δ vanishes on an open set of times). So we can assume that the SLEs stay in D_1, D_2 , hence we have uniformity in the $O(n^{-3})$ estimate of the commutation condition. The last case to study is when the trace gets close to the boundary, say $|x_{i-1} - z_j| < 2\delta$ for some i, j , without actually crossing it. The probability of this event goes to zero as N goes to infinity and δ goes to zero.

So the above estimate is valid up to an event of negligible probability, viz. either an SLE crosses ∂D_1 or ∂D_2 without the functional vanishing or one of the f_δ is less than one and yet the functional does not vanish. Taking the limit as n goes to infinity and $N \nearrow \infty$, $\delta \searrow 0$ (so that $C(N, \delta)/n \rightarrow 0$), one gets the stated identity, that is :

$$\mathbb{E}_1(\mathbf{1}_{\tau_1 > S, \tau_2 > T}) = \mathbb{E}_2(\mathbf{1}_{\tau_1 > S, \tau_2 > T}).$$

This concludes the proof of the lemma. □

Proof of the Proposition. Let D_1, D_2 be as in the lemma. We grow an $\text{SLE}_\kappa(b)$ in D_1 until it reaches ∂D_1 , and then in the remaining domain an $\text{SLE}_{\tilde{\kappa}}(\tilde{b})$ in D_2 until it reaches ∂D_2 . This defines a Loewner chain $(K_{s,t})_{(s,t) \in \mathcal{T}}$. We will prove that this chain is an $\text{SLE}(\kappa, b, \tilde{\kappa}, \tilde{b})$, i.e. it has the appropriate Markov property.

Let $0 = S_0 < S_1 < \dots < S_k = \infty$ and $0 = T_0 < T_1 < \dots < T_k = \infty$ be sequences of fixed times. Let σ be a permutation of the symbols $(S_1, \dots, S_k, T_1, \dots, T_k)$, which is increasing for the partial order generated by $S_i < S_{i+1}, T_i < T_{i+1}$. Let $(K_{s,t}^\sigma)$ be the (random) Loewner chain obtained by growing SLEs alternatively in D_1 and in D_2 according to σ , stopping the SLEs when they reach $\partial D_1, \partial D_2$. For instance, if $\sigma = (S_1, T_1, T_2, S_2, \dots)$, one grows the first SLE to half-plane capacity $2S_1$ (and stop it if it reaches ∂D_1), then the second SLE to half-plane capacity $2S_2$, measured in the original half-plane (and stop it if it reaches ∂D_1), and then again the first SLE to half-plane capacity $2S_2$ (and stop it ...), ... To alleviate notations, we will use the convention that for a Loewner chain $(\tilde{K}_{s,t})$, $\tilde{K}_{s,t} = \tilde{K}_{s \wedge S, t \wedge T}$, where S is the time at which $(\tilde{K}_{s,0})_s$ exits D_1 (resp. T is the time at which $(\tilde{K}_{0,t})_t$ exits D_2).

If σ and σ' differ by a single transposition, i.e. $\sigma = (\sigma_1, S_{k_1}, T_{k_2}, \sigma_2)$, $\sigma' = (\sigma_1, T_{k_2}, S_{k_1}, \sigma_2)$, then we can apply the lemma with $\phi_0 = \phi_{K_{S_{k_1}, T_{k_2}}^\sigma}$, $D'_1 = \phi_0(D_1)$, $D'_2 = \phi_0(D_2)$. This proves that we can couple $(K_{s,t}^\sigma)$

and $(K_{s,t}^{\sigma'})$. By induction, we can couple (simultaneously) the chains $(K_{s,t}^{\sigma})$ for all admissible permutations σ . By construction, for $\sigma = (S_1, \dots, S_k, T_1, \dots, T_k)$, K^{σ} is distributed as K .

Now, for any $k_1, k_2 \in \{0, \dots, k\}$, one can consider the permutation:

$$\sigma = (S_0, \dots, S_{k_1}, T_0, \dots, T_{k_2}, S_{k_1+1}, \dots, S_k, T_{k_2+1}, \dots, T_k).$$

The previous coupling proves the Markov property for the fixed time (S_{k_1}, T_{k_2}) (i.e the chain

$$(g_{S_{k_1}, T_{k_2}}(K_{S_1+s, T_2} \setminus K_{S_1, T_2}))$$

is a stopped $\text{SLE}_{\kappa}(b)$, and the same thing holds for the other SLE).

This still holds for stopping times supported on $\{(S_{k_1}, T_{k_2}), k_1, k_2 = 0, \dots, k\}$. Since the subdivisions $S_0 < \dots < S_k$ and $T_0 < \dots < T_k$ were arbitrary, this also holds for stopping times with finite support, and by a limiting argument for all stopping times (as for the classical Markov property).

□

7 Classification of commuting SLEs

We can now conclude the general study of pairs of commuting chordal SLEs in a simply connected domain. In the upper half-plane $\mathbb{H}$, with $(2n+2)$ marked points $(x, y, z_1, \dots, z_n)$ on the real line (and one marked point z_{n+1} at infinity), consider two parameters $\kappa, \tilde{\kappa}$, and two smooth functions of the configuration $(x, y, z_1, \dots, z_n)$, translation invariant and homogeneous of degree -1 . Let $\mathcal{L}$ be the infinitesimal generator of the $\text{SLE}_{\kappa}(b)$ growing at x (driven by (X_s) , (g_s) are the corresponding conformal equivalences), and $\mathcal{M}$ the infinitesimal generator of the $\text{SLE}_{\tilde{\kappa}}(\tilde{b})$ growing at y (driven by (Y_t) , $(\tilde{g}_t)$ are the corresponding conformal equivalences). By a cocycle $C(\phi, z)$ we mean a function of the form:

$$C(\phi, z) = \prod_{1 \leq i < j \leq n+1} \left(\phi'(z_i) \phi'(z_j) \left(\frac{z_j - z_i}{\phi(z_j) - \phi(z_i)} \right)^2 \right)^{\nu_{ij}} \frac{f(\phi(z_1), \dots, \phi(z_n))}{f(z_1, \dots, z_n)}$$

where f is a non-vanishing function (translation invariant and homogeneous of degree 0). So f is a conformally invariant function of the marked points $(z-1, \dots, z_n, z_{n+1} = \infty)$ and can be seen as a function on the residual (i.e. not involving the positions of x, y) moduli space. If $n \geq 3$, this decomposition is not unique, as discussed in Section 4.

Theorem 7. *The following assertions are equivalent.*

(i). *There exists an $\text{SLE}(\kappa, b, \tilde{\kappa}, \tilde{b})$.*

(ii). *The infinitesimal generators satisfy the relation:*

$$[\mathcal{L}, \mathcal{M}] = \frac{4}{(y-x)^2} (\mathcal{M} - \mathcal{L})$$

(iii). *$\tilde{\kappa} = \kappa$ or $\tilde{\kappa} = 16/\kappa$, $b = \kappa \partial_x \psi / \psi$, $\tilde{b} = \tilde{\kappa} \partial_y \psi / \psi$, where ψ is a non-vanishing solution of the system:*

$$\begin{cases} \frac{\kappa}{2} \partial_{xx} \psi + \sum_i \frac{2\partial_i \psi}{z_i - x} + \frac{2\partial_y \psi}{y - x} + \left(\left(1 - \frac{6}{\tilde{\kappa}}\right) \frac{1}{(y-x)^2} + h(x, z) \right) \psi = 0 \\ \frac{\tilde{\kappa}}{2} \partial_{yy} \psi + \sum_i \frac{2\partial_i \psi}{z_i - y} + \frac{2\partial_x \psi}{x - y} + \left(\left(1 - \frac{6}{\kappa}\right) \frac{1}{(x-y)^2} + h(y, z) \right) \psi = 0 \end{cases}$$

where

$$h(x, z) = \sum_i \frac{\mu_i}{(x - z_i)^2} + \sum_{i < j} \frac{\nu_{ij}}{(x - z_i)(x - z_j)} + \ell_x \log(f(z)),$$

the μ_i, ν_{ij} are constant parameters, and f is a function on the residual moduli space.

(iv). $\tilde{\kappa} = \kappa$ or $\tilde{\kappa} = 16/\kappa$, and there is a non-vanishing function ψ and a cocycle C such that if:

$$\begin{aligned} Z_s &= \psi(X_s, g_s(y), \dots, g_s(z_i), \dots) g'_s(y)^{\alpha_{\tilde{\kappa}}} C(g_s, z) \\ \tilde{Z}_t &= \psi(\tilde{g}_t(x), Y_t, \dots, \tilde{g}_t(z_i), \dots) \tilde{g}'_t(x)^{\alpha_{\tilde{\kappa}}} C(\tilde{g}_t, z) \end{aligned}$$

then (Z_s) is a local martingale for chordal $\text{SLE}_{\kappa}(x \rightarrow \infty)$, $(\tilde{Z}_t)$ is a local martingale for chordal $\text{SLE}_{\tilde{\kappa}}(y \rightarrow \infty)$, and $\text{SLE}_{\kappa}(b)$ is the Girsanov transform of chordal SLE_{κ} by (Z_s) , $\text{SLE}_{\tilde{\kappa}}(\tilde{b})$ is the Girsanov transform of chordal $\text{SLE}_{\tilde{\kappa}}$ by $(\tilde{Z}_t)$.

Note that there is no loss of generality in considering two (rather than $m \geq 2$) commuting SLEs. Indeed, the only conditions will be the pairwise conditions. It is also easy to see that the proofs for local commutation can be adapted for m SLEs (though notations become quite heavy). Let us explicit, say, condition (iii) in this situation. On the real line, $(m+n)$ points $(y_1, \dots, y_m, z_1, \dots, z_n)$ are marked, and we want to grow m SLEs $(\text{SLE}_{\kappa_i}(b_i), i = 1, \dots, m)$ at $y_1, \dots, y_m$. Then $\{\kappa_1, \dots, \kappa_m\} \subset \{\kappa_1, 16/\kappa_1\}$, $b_i = \kappa_i \partial_{y_i} \psi / \psi$, where ψ is annihilated by the operators:

$$\frac{\kappa}{2} \partial_{y_i y_i} + \sum_{j \neq i} \frac{2 \partial_{y_j}}{y_j - y_i} + \sum_j \frac{2 \partial_{z_j}}{z_j - y_i} - 2 \left(\sum_{j \neq i} \frac{\alpha_{\kappa_j}}{(y_j - y_i)^2} + \sum_j \frac{\mu'_j}{(z_j - y_i)^2} + \sum_{j < j'} \frac{\nu'_{jj'}}{(z_j - y_i)(z_{j'} - y_i)} + \ell_{y_i} \log(f(z)) \right)$$

for some parameters $\mu'_j, \nu'_{jj'}$ and some function f on the residual moduli space.

8 Restriction formulae for non-intersecting SLEs

In this section we specialize to a simple parametric case, where n SLEs started from distinct points on the real line are aiming at infinity; there are only n marked points on the real line (and one at infinity). Each of the n SLEs is an $\text{SLE}_{\kappa}(\underline{\rho})$, where $\underline{\rho} = (2, \dots, 2)$. In this situation, we can not only define locally a n -parameter Loewner chain, but also define it globally if $\kappa \leq 4$. Indeed, the only thing preventing from a global definition is the possibility of collisions of marked points. But such collisions a.s. don't happen for these $\text{SLE}_{\kappa}(2, \dots, 2)$, so we can actually define a chain with full time set $\mathbb{R}_+^n$.

If the n starting points collapse to zero, we get n “non-intersecting” SLEs starting at 0 and ending at ∞ . Restriction formulae are derived for these Loewner chains (indexed by $\mathbb{R}_+^n$). This gives a simple realization of the exponents $h_{1;n+1}(\kappa)$ (see also [23]).

The radial case (n “non-intersecting” SLEs started from the boundary and aiming at a single bulk point) is also studied, and restriction formulae then give the exponent $2h_{0;n/2}(\kappa)$.

8.1 The chordal case

Let $y_1 < \dots < y_n$ be n real points. Consider the infinitesimal generators:

$$\mathcal{L}_i = \frac{\kappa}{2} \partial_{ii} + \left(\sum_{j \neq i} \frac{2}{y_i - y_j} \right) \partial_i + \sum_{j \neq i} \frac{2}{y_j - y_i} \partial_j$$

Then, from the previous computations (parametric case), we see that the following commutation relations are satisfied:

$$[\mathcal{L}_i, \mathcal{L}_j] = \frac{4}{(y_j - y_i)^2} (\mathcal{L}_j - \mathcal{L}_i).$$

As mentioned earlier, this ensures (if $\kappa \leq 4$) the existence of a process $(K_{s_1, \dots, s_n})$ such that: $(\phi_{K_{s_1, \dots, s_n}}(K_{s_1, \dots, s_i + s, \dots, s_n}))_{s \geq 0}$ is an $\text{SLE}_\kappa(2, \dots, 2)$ started from $(\phi(\gamma_{s_i}^i)) = (\phi(\gamma_{s_1}^1), \dots, \phi(\gamma_{s_n}^n))$, independent from $(K_{t_1, \dots, t_n})_{t_j \leq s_j}$, where $\phi = \phi_{K_{s_1, \dots, s_n}}$, $K_{0, \dots, s_i, \dots, 0} = \gamma_{[0, s_i]}^i$. Assume that this process is started from $(y_1, \dots, y_n)$, i.e. $\gamma_0^i = y_i$, and define $K_\infty = \bigcup_i \gamma_{[0, \infty]}^i \subset \overline{\mathbb{H}}$.

Consider now a hull $A \subset \overline{\mathbb{H}}$, that does not intersect $\{y_1, \dots, y_n\}$. Let $\lambda_\kappa = (6 - \kappa)(8 - 3\kappa)/2\kappa$. If $\kappa \leq 8/3$, and L is an independent random loop soup with intensity λ_κ in $\mathbb{H}$, define K_∞^L to be (the filling of) the union of K_∞ and the loops in L that intersect it. Then, if $\kappa \leq 8/3$, the following restriction formula holds:

Lemma 8. *The probability that K_∞^L does not intersect A is given by:*

$$\mathbb{P}(K_\infty^L \cap A = \emptyset) = \prod_i \phi'_A(y_i)^{(6-\kappa)/2\kappa} \prod_{i < j} \left(\frac{\phi_A(y_j) - \phi_A(y_i)}{y_j - y_i} \right)^{2/\kappa}$$

Proof. Define $\underline{s} = (s_1, \dots, s_n)$, $h_{\underline{s}} = \phi_{\phi_{K_{\underline{s}}}(A)}$, and $Y_{\underline{s}}^{(i)} = \phi_{\underline{s}}(\gamma_{s_i}^i)$. Then, from the definition of $K_{\underline{s}}$ and Lemma 4 in [7], one sees that:

$$s_k \mapsto \prod_i h'_{\underline{s}}(Y_{\underline{s}}^{(i)})^{(6-\kappa)/2\kappa} \prod_{i < j} \left(\frac{h_{\underline{s}}(y_j) - h_{\underline{s}}(y_i)}{y_j - y_i} \right)^{2/\kappa} \exp \left(\lambda_\kappa \int_0^{s_k} \frac{S h_{s_1, \dots, s_k, \dots, s_n}(Y_{s_1, \dots, s_k, \dots, s_n}^{(k)})}{6} d\sigma_k \right)$$

is a bounded martingale for all $k, s_1, \dots, s_n$. From the properties of the Brownian loop soup, it appears that the following semimartingale (proportional to the first one) is also a bounded martingale:

$$s_k \mapsto \prod_i h'_{\underline{s}}(Y_{\underline{s}}^{(i)})^{(6-\kappa)/2\kappa} \prod_{i < j} \left(\frac{h_{\underline{s}}(y_j) - h_{\underline{s}}(y_i)}{y_j - y_i} \right)^{2/\kappa} \mathbb{P}(K_{\underline{s}}^L \cap A = \emptyset)$$

Hence, for all $\underline{s} = (s_1, \dots, s_n)$, one gets (using n different martingales):

$$\prod_i \phi'_A(y_i)^{(6-\kappa)/2\kappa} \prod_{i < j} \left(\frac{\phi_A(y_j) - \phi_A(y_i)}{y_j - y_i} \right)^{2/\kappa} = \mathbb{E} \left(\prod_i h'_{\underline{s}}(Y_{\underline{s}}^{(i)})^{(6-\kappa)/2\kappa} \prod_{i < j} \left(\frac{h_{\underline{s}}(y_j) - h_{\underline{s}}(y_i)}{y_j - y_i} \right)^{2/\kappa} \mathbb{P}(K_{\underline{s}}^L \cap A = \emptyset) \right)$$

Now, as $\inf \underline{s}$ goes to infinity, the product in the right-hand side converges to $\mathbb{P}(K_\infty^L \cap A = \emptyset)$, which concludes the proof. $\square$

Define the conformal weight $h_{p;q} = h_{p;q}(\kappa)$ by:

$$h_{p;q} = \frac{(p\kappa - 4q)^2 - (\kappa - 4)^2}{16\kappa}$$

Then, if $y_1, \dots, y_n$ collapse to zero, the above formula reduces to:

$$\mathbb{P}(K_\infty^L \cap A = \emptyset) = \phi'_A(0)^{h_{1;n+1}}$$

The role of the conformal weights $h_{1;n+1}$ in the context of restriction measures and $\text{SLE}_\kappa(\rho)$ is discussed in [23].

Corollary 9 (Restriction property). *Let $\kappa \leq 8/3$, L a loop soup with intensity λ_κ . Conditionally on $\{K_\infty^L \cap A = \emptyset\}$ and up to a time change, $((\phi_A(K_{\underline{s}}))_{\underline{s}}, \phi_A(\{\delta \in L : \delta \cap A = \emptyset\}))$ is distributed as $((K_{\underline{s}})_{\underline{s}}, L)$, where $(K_{\underline{s}})$ is started from $\phi_A(y_1), \dots, \phi_A(y_n)$. In particular, the collection of measures on hulls K_∞^L indexed by the starting configuration $(\mathbb{H}, z_1, \dots, z_n, z_{n+1} = \infty)$ has the restriction property.*

Proof. From the previous lemma, the result is a straightforward application of the Girsanov theorem and the restriction property of the loop soup. The assertion on the restriction property can be derived directly by applying the previous formula to concatenation of hulls $A.B$. $\square$

8.2 The radial case

Recall the definition of radial $\text{SLE}_\kappa(\underline{\rho})$: assume the existence of processes $(\xi_t)_{t \geq 0}$ and $(\chi_t^{(i)})_{t \geq 0}$, $i \in \{1 \dots k\}$, satisfying the SDEs:

$$\begin{cases} d\xi_t = (i\xi_t \sqrt{\kappa} dB_t - \frac{\kappa}{2} \xi_t dt) + \sum_{i=1}^k \frac{\rho_i}{2} \left(-\xi_t \frac{\xi_t + \chi_t^{(i)}}{\xi_t - \chi_t^{(i)}} \right) dt \\ d\chi_t^{(i)} = -\chi_t \frac{\chi_t + \xi_t}{\chi_t - \xi_t} dt \end{cases}$$

Then the ODEs $dg_t(z) = -g_t(z)(g_t(z) + \xi_t)/(g_t(z) - \xi_t)dt$ define radial $\text{SLE}_\kappa(\underline{\rho})$ in the unit disk $\mathbb{U}$.

First, we briefly discuss commutation conditions in the radial case. Suppose that $\chi_1, \dots, \chi_n$ are n points on the unit circle. One considers two SLEs growing at χ_1 and χ_n resp., assuming that the drift terms are functions of the χ_i . We think of functions annihilated by infinitesimal generators as expected values of some events; it is quite natural to express these real-valued functions in angular coordinates: $\chi_j = \exp(i\theta_j)$. Reasoning as in the chordal case, if $\mathcal{L}$ and $\mathcal{M}$ are the infinitesimal generators of the two SLEs, the commutation condition reads:

$$[\mathcal{L}, \mathcal{M}] = \frac{1}{\sin((\theta_n - \theta_1)/2)^2} (\mathcal{L} - \mathcal{M}).$$

The generator for a radial $\text{SLE}_\kappa(\rho_1, \dots, \rho_{n-1})$ started from $(\chi_1, \dots, \chi_n)$ is:

$$\mathcal{L} = \frac{\kappa}{2} \partial_{11} + \sum_{i>1} \cot\left(\frac{\theta_i - \theta_1}{2}\right) \left(\partial_i - \frac{\rho_i}{2} \partial_1\right)$$

By analogy with the chordal case, one can find solutions for this commutation relation:

- (i). Two $\text{SLE}_\kappa(\rho)$ started from (χ_1, χ_2) and (χ_2, χ_1) resp., $\rho \in \{2, \kappa - 6\}$.
- (ii). n $\text{SLE}_\kappa(2, \dots, 2)$ started from $(\chi_i, \chi_1, \dots, \widehat{\chi_i}, \dots, \chi_n)$.

Let us comment briefly on the case (i). For $\rho = \kappa - 6$, this is only chordal reversibility in a radial normalization (as may be seen by slightly modifying the argument for chordal-radial equivalence when $\kappa = 6$). In the case $\rho = 2$, this gives a model of “pinned chordal SLE”, i.e chordal SLE “conditionally” on the trace visiting a given bulk point, for $\kappa < 8$. More precisely, start from a chordal SLE in radial normalization (hence, up to a time change, radial $\text{SLE}_\kappa(\kappa - 6)$). Then the first moment estimate in [3] relies on the computation of the leading eigenvector for the associated infinitesimal generator. This yields a local martingale:

$$s \mapsto \sin\left(\frac{\theta_s^{(1)} - \theta_s^{(2)}}{2}\right)^{8/\kappa - 1} e^{(1 - \kappa/8)s}$$

corresponding of the probability that the SLE trace gets infinitely close to the bulk point 0. Using this as a Girsanov density, one gets a radial $\text{SLE}_\kappa(2)$. Note that for $\kappa = 8$, this density is 1, and $\kappa - 6 = 2$.

There are other examples with two boundary points. Consider a chordal SLE_κ from χ_1 to χ_2 and condition it to leave 0 on its left (resp. right); this can be made explicit (see [20]). Once again, the drift terms are (generically in κ) transcendental.

In the case (ii), just as in the chordal case (at least if $\kappa \leq 4$), based on the infinitesimal commutation relations, one can define a n -braids radial SLE. The question of such a definition, from a CFT point of view, appears in [6, 5]. Note also that summing the n generators here gives the generator of Dyson's Brownian motion.

As above, we study the case (ii) from the restriction point of view. First, we have to derive restriction formulae for radial $\text{SLE}_\kappa(\rho)$. Let A be a hull of $\overline{\mathbb{U}}$ (i.e. A is a compact subset of $\overline{\mathbb{U}}$, $A \cap \overline{\mathbb{U}} = \overline{A \cap \mathbb{U}}$, $\mathbb{U} \setminus A$ is equivalent to $\mathbb{U}$ and $0 \notin A$). For any such hull, denote by ϕ_A the unique conformal equivalence $\mathbb{U} \setminus A \rightarrow \mathbb{U}$ such that $\phi_A(0) = 0$ and $\phi'_A(0) > 0$. Suppose that $\xi_0, \chi_0^{(i)}$ are not in A . Then $h_t = \phi_{g_t(A)}$ is defined at least for small times. Recall that $\lambda_\kappa = (8 - 3\kappa)(6 - \kappa)/2\kappa$. Then the following result (analogous to Lemma 4 in [7] and generalizing a result stated in [13]) holds:

Lemma 10. *In this situation, define $\tilde{\xi}_t = h_t(\xi_t)$, $\tilde{\chi}_t^{(i)} = h_t(\chi_t^{(i)})$:*

$$\begin{aligned} M_t^\varnothing &= \left(\frac{h'_t(\xi_t)\xi_t}{\tilde{\xi}_t} \right)^{\frac{(6-\kappa)}{2\kappa}} h'_t(0)^{\frac{(6-\kappa)(\kappa-2)}{8\kappa}} \exp \left(-\lambda_\kappa \int_0^t \frac{\xi_s^2 S h_s(\xi_s)}{6} ds \right) \\ M_t^{(i)} &= \left(\frac{h'_t(\chi_t^{(i)})\chi_t^{(i)}}{\tilde{\chi}_t^{(i)}} \right)^{\frac{\rho_i(\rho_i+4-\kappa)}{4\kappa}} \left(\frac{(\tilde{\chi}_t^{(i)} - \tilde{\xi}_t)^2}{\tilde{\chi}_t^{(i)}\tilde{\xi}_t} \frac{\chi_t^{(i)}\xi_t}{(\chi_t^{(i)} - \xi_t)^2} \right)^{\frac{\rho_i}{2\kappa}} h'_t(0)^{\frac{\rho_i(\rho_i+4)}{8\kappa}} \\ M_t^{(i,j)} &= \left(\frac{(\tilde{\chi}_t^{(i)} - \tilde{\chi}_t^{(j)})^2}{\tilde{\chi}_t^{(i)}\tilde{\chi}_t^{(j)}} \frac{\chi_t^{(i)}\chi_t^{(j)}}{(\chi_t^{(i)} - \chi_t^{(j)})^2} \right)^{\frac{\rho_i\rho_j}{4\kappa}} h'_t(0)^{\frac{\rho_i\rho_j}{4\kappa}} \end{aligned}$$

Note that all fractions are real numbers. Then $(M^\varnothing \prod_i M^{(i)} \prod_{i < j} M^{(i,j)})$ is defined up to some random positive time (possibly infinite), and is a local martingale.

Proof. This is a rather straightforward transposition to the radial case of results and methods in [13], which we discuss for the sake of completeness. We use freely a complex-variable version of Itô's formula.

Let $\tilde{g}_t = \phi_{\phi_A(K_t)}$, with the usual notations, so that $h_t \circ g_t = \tilde{g}_t \circ \phi_A$. Then $(\tilde{g}_t)$ is a time-changed radial Loewner chain:

$$\partial_t \tilde{g}_t = -\tilde{g}_t \frac{\tilde{g}_t + \tilde{\xi}_t}{\tilde{g}_t - \tilde{\xi}_t} da_t$$

and $da_t = (h'_t(\xi_t)\xi_t/\tilde{\xi}_t)^2$. This is the key “commutative diagram” argument of [13]. Then, standard differential calculus yields:

$$\begin{aligned} \partial_t h_t &= -h_t \frac{h_t + \tilde{\xi}_t}{h_t - \tilde{\xi}_t} \partial_t a_t + h'_t \cdot z \frac{z + \xi_t}{z - \xi_t} \\ [\partial_t h_t](\xi_t) &= 3\xi_t h'_t(\xi_t) + 3\xi_t^2 h''_t(\xi_t) - 3 \frac{(\xi_t h'_t(\xi_t))^2}{\tilde{\xi}_t} \\ \partial_t h'_t &= -h'_t \frac{h_t + \tilde{\xi}_t}{h_t - \tilde{\xi}_t} \partial_t a_t + h_t h'_t \frac{2\tilde{\xi}_t}{(h_t - \tilde{\xi}_t)^2} \partial_t a_t + h''_t \cdot z \frac{z + \xi_t}{z - \xi_t} + h'_t \cdot \left(\frac{z + \xi_t}{z - \xi_t} - \frac{2z\xi_t}{(z - \xi_t)^2} \right) \\ [\partial_t h'_t](\xi_t) &= \left(\frac{4}{3} \xi_t^2 h'''_t - \frac{(\xi_t h''_t)^2}{2h'_t} + 3\xi_t h''_t - \frac{\xi_t^2 (h'_t)^3}{\tilde{\xi}_t^2} + h'_t \right) (\xi_t) \\ \partial_t h'_t(0) &= h'_t(0)(\partial_t a_t - 1) \end{aligned}$$

Relax the assumptions for the moment, and assume that ξ satisfies the SDE:

$$d\xi_t = i\xi_t\sqrt{\kappa}dB_t - \frac{\kappa}{2}\xi_t dt + \xi_t b_t dt$$

where B is a standard Brownian motion and b_t is a (progressive) drift coefficient. Define:

$$\begin{aligned}\sigma_t &= \frac{\xi_t h_t''(\xi_t)}{h_t'(\xi_t)} + 1 - \frac{\xi_t h_t'(\xi_t)}{\tilde{\xi}_t} \\ \sigma_t^{(i)} &= \frac{\chi_t^{(i)} + \xi_t}{\chi_t^{(i)} - \xi_t} - \frac{\tilde{\chi}_t^{(i)} + \tilde{\xi}_t}{\tilde{\chi}_t^{(i)} - \tilde{\xi}_t} \frac{\xi_t h_t'(\xi_t)}{\tilde{\xi}_t}\end{aligned}$$

Then:

$$\begin{aligned}\frac{dM_t^\varnothing}{M_t^\varnothing} &= \frac{6-\kappa}{2\kappa}\sigma_t (i\sqrt{\kappa}dB_t + b_t dt) \\ \left(\frac{\rho_i}{2\kappa}\right)^{-1} \frac{dM_t^{(i)}}{M_t^{(i)}} &= \sigma_t^{(i)} \left(i\sqrt{\kappa}dB_t + b_t dt + \frac{\rho_i}{2} \frac{\xi_t + \chi_t^{(i)}}{\xi_t - \chi_t^{(i)}} dt \right) - \frac{(6-\kappa)}{2}\sigma_t \frac{\tilde{\chi}_t^{(i)} + \tilde{\xi}_t}{\tilde{\chi}_t^{(i)} - \tilde{\xi}_t} \frac{\xi_t h_t'(\xi_t)}{\tilde{\xi}_t} dt \\ \left(\frac{\rho_i \rho_j}{4\kappa}\right)^{-1} \frac{\partial_t M_t^{(i,j)}}{M_t^{(i,j)}} &= \sigma_t^{(i)} \sigma_t^{(j)} + \frac{\xi_t + \chi_t^{(i)}}{\xi_t - \chi_t^{(i)}} \sigma_t^{(j)} + \frac{\xi_t + \chi_t^{(j)}}{\xi_t - \chi_t^{(j)}} \sigma_t^{(i)}\end{aligned}$$

Given that $b_t = -\sum_i \frac{\rho_i}{2} \frac{\xi_t + \chi_t^{(i)}}{\xi_t - \chi_t^{(i)}}$, if N denotes $(M^\varnothing \prod_i M^{(i)} \prod_{i<j} M^{(i,j)})$, one gets:

$$\begin{aligned}\frac{dN_t}{N_t} &= \frac{dM_t^\varnothing}{M_t^\varnothing} + \sum_i \left(\frac{dM_t^{(i)}}{M_t^{(i)}} + \frac{d\langle M_t^\varnothing, M_t^{(i)} \rangle}{M_t^\varnothing M_t^{(i)}} \right) + \sum_{i<j} \left(\frac{dM_t^{(i,j)}}{M_t^{(i,j)}} + \frac{d\langle M_t^{(i)}, M_t^{(j)} \rangle}{M_t^{(i)} M_t^{(j)}} \right) \\ &= \sigma_t' dB_t + \frac{6-\kappa}{2\kappa} b_t \sigma_t + \sum_i \left(\frac{6-\kappa}{2\kappa} \sigma_t \frac{\rho_i}{2} \frac{\xi_t + \chi_t^{(i)}}{\xi_t - \chi_t^{(i)}} + \frac{\rho_i}{2\kappa} \sigma_t^{(i)} \left(b_t + \frac{\rho_i}{2} \frac{\xi_t + \chi_t^{(i)}}{\xi_t - \chi_t^{(i)}} \right) \right) dt \\ &\quad + \sum_{i<j} \frac{\rho_i \rho_j}{4\kappa} \left(\frac{\xi_t + \chi_t^{(i)}}{\xi_t - \chi_t^{(i)}} \sigma_t^{(j)} + \frac{\xi_t + \chi_t^{(j)}}{\xi_t - \chi_t^{(j)}} \sigma_t^{(i)} \right) dt \\ &= \sigma_t' dB_t\end{aligned}$$

□

As in the chordal case, the symmetry of these formulae when $\underline{\rho} = (2, \dots, 2)$ enables to derive restriction formulae for n -braid SLEs.

Lemma 11. *Let $(K_\cdot)$ be a radial n -braid SLE in $\mathbb{U}$ started from distinct points $\chi_1, \dots, \chi_n$, and A be a hull not intersecting these points. If $\kappa \leq 8/3$, and L is an independent loop soup with intensity λ_κ , then:*

$$\mathbb{P}(K_\infty^L \cap A = \varnothing) = |\phi_A'(0)|^{2h_{0;n/2}} \prod_i |\phi_A'(\chi_i)|^{h_{1;2}} \prod_{i<j} \left| \frac{\phi_A(\chi_j) - \phi_A(\chi_i)}{\chi_j - \chi_i} \right|^{2/\kappa}$$

If $\chi_0^{(1)}, \dots, \chi_0^{(n)}$ collapse to χ , the above formula reduces to:

$$|\phi_A'(\chi)|^{h_{1;n+1}} |\phi_A'(0)|^{2h_{0;n/2}}$$

Corollary 12 (Restriction property). *Let $\chi_1, \dots, \chi_n$ and K_∞^L as above. This defines a family of probability measures $\mu_{\kappa, \underline{\chi}}$ on “sea star” hulls $K' = K^L$. This family has the restriction property: for all hull A , $(\phi_A)_* \mu_{\kappa, \underline{\chi}}(\cdot | K' \cap A = \varnothing) = \mu_{\kappa, (\phi_A)_*(\underline{\chi})}$*

Proof. As in the chordal case. □

9 Multiply connected domains

One can think of SLE as a diffusion in a configuration space. The diffusion coefficients are constrained by the conformal invariance requirement. If the associated moduli space is a point, then the coefficients are constant parameters; this situation corresponds to chordal and radial SLE, and $\text{SLE}(\kappa, \rho)$ (and also “annulus SLE”). If the moduli space is larger, then SLE is essentially specified by the data of diffusion coefficients as functions on the moduli space; so we are no longer in the parametric situation. We only discuss the “constant κ ” case, for physical and technical reasons. Also, we will be mainly interested in expressing necessary conditions for reversibility in multiply connected domains, so we will not carry the discussion in the same degree of generality as in the simply connected case.

In the case of a simply connected domains with $2n$ points marked on the boundary, SLE is specified by κ and a function of $(2n - 3)$ independent cross-ratios of the $2n$ boundary points. If we add the requirement that the SLE commutes with $(2n - 1)$ SLEs started at the other points, then we have to choose a “partition function” ψ as discussed earlier. This function belongs to the finite-dimensional solution space of a holonomic system derived from the commutation conditions, and the situation is parametric again ($1 + C(2n, n)$ parameters).

Similarly, in the case of multiply connected domain, we want to restrict the diffusion coefficients to the “physically relevant” ones. We consider in particular the case of chordal SLE (going from x to y , x and y on the same boundary component) in a multiply connected domain.

There are at least two ways to describe SLE in a multiply connected domain. The first one, that follows closely the simply connected case, consists in choosing a parametric family of standard domains (a section of the moduli space), and writing explicit diffusion equations for the parameters; this is the approach of [8, 2]. Another route, following Makarov and Zhan (see [27]), consists in using a local chart at the growth point and a “conformally invariant SDE”, so that the path distribution does not depend on the choice of local chart. In the first case, the diffusion coefficient is a function on the moduli space; in the other case, SLE is specified by a “partition function”, which is a conformally covariant function on the configuration space; taking its log derivative (w.r.t. the growth point), one gets a function on the moduli space. We will use this second framework, that better suits our purposes.

So let $\mathfrak{C}$ be the configuration space of $(g + 1)$ -connected plane domains with $(m + 2)$ points marked on the boundary and n points marked in the bulk. Two of the marked points, x and y are on the same component of the boundary. Denote by $\mathfrak{M}$ the associated moduli space.

First we briefly summarize the local chart approach. Any configuration is equivalent to a configuration of type $(\mathbb{H} \setminus K, x, \dots)$ where x is real and K is a compact subset of $\mathbb{H}$ (with g connected component). By conformal invariance, we need only to define SLE for these configurations, and need to do it coherently (independently of choices). Let h be a conformal equivalence between $c = (\mathbb{H} \setminus K, x, y = \infty, \dots)$ and $h_*c = (\mathbb{H} \setminus K', x', y' = \infty, \dots)$. SLE in c is defined by the chordal Loewner equations and an SDE:

$$\partial_t g_t = \frac{2}{g_t - W_t}, dW_t = \sqrt{\kappa} dB_t + b(c_t) dt$$

From [13], we can write the SDE for the driving process of the image of the SLE by $h_0 = h$:

$$dh_t(W_t) = h'_t(W_t) dW_t + \left(\frac{\kappa}{2} - 3\right) h''_t(W_t) dt$$

where $h_t = \tilde{g}_t \circ h \circ g_t^{-1}$, and $\tilde{g}_t$ also solves the chordal Loewner equations (though with a time change). After a time change, and at time 0, one sees that the condition necessary for invariance of the SDE is the following covariance condition:

$$b(h_*c) = \frac{b(c)}{h'(x)} + \left(\frac{\kappa}{2} - 3\right) \frac{h''}{h'^2}(x).$$

Here h is normalized by $h(\infty) = \infty$, $h'(\infty) = 1$ (hydrodynamic normalization at infinity). Let ψ be a positive function on the configuration space. We say that ψ is α -covariant if:

- (i). (Möbius invariance) For any $c = (D \setminus K, \dots)$, $c' = (D' \setminus K', \dots)$ where D, D' are simply connected, $h : c \rightarrow c' = h_*c$ is an equivalence of configurations, and h extends to a conformal equivalence $D \rightarrow D'$, one has:

$$\psi(h_*c) = \psi(c)$$

- (ii). (covariance) For any $c = (D \setminus K, x, y, \dots)$, $c' = (D \setminus K', x, y, \dots)$ where D is simply connected, $x, y \in \partial D$, ∂D is smooth at x, y , and $h : c \rightarrow c' = h_*c$ is an equivalence of configurations, one has:

$$\psi(h_*c) = (h'(x)h'(y))^{-\alpha}\psi(c)$$

Note that the function ψ is completely determined by these conditions and its restriction to a section of the moduli space. Let us give three (important) examples of such covariant functions, say for annuli with two marked points on one component of the boundary: $c = (D, x, y)$.

- (i). In $c = (D, x, y)$, assume that the arc (xy) is blue and (yx) is yellow. Let $\psi(c)$ be the probability that (xy) and (yx) are connected to the other boundary component by a blue (resp. yellow) cluster in the scaling limit of critical percolation. Alternatively, $\psi(c)$ is the corresponding SLE_6 probability (see [8]). Then ψ is 0-covariant. (Also, $\psi = 1$ is 0-covariant; this is a version of locality for SLE_6).
- (ii). In his thesis, Beffara uses the results of [13] and an inclusion-exclusion argument to prove the following: let $\psi(c)$ be the probability that chordal $\text{SLE}_{8/3}$ from x to y in the (filled) domain avoids the hole (resp. leaves the hole on its left, resp. leaves the hole on its right). Then ψ is 5/8-covariant.
- (iii). If $\kappa = 2$, $\alpha_\kappa = 1$, and SLE_2 is the scaling limit of Loop-Erased Random Walks ([16, 26]); these walks are closely related to some discrete harmonic quantities. Define:

$$\psi(c) = \frac{\partial^2}{\partial n_x \partial n_y} G_D(w, w')$$

the normal derivative at x and y of the Green kernel (which is symmetric in the two variables). Then the invariance property of the Green kernel implies that ψ is 1-covariant. In general domains, this is the (chordal version of) Harmonic Random Loewner Chain (HRLC) as defined by Zhan in [27]. Similar harmonic constructions exist for $\kappa = 8$, $\alpha_\kappa = -1/8$, using normal reflection on some boundary components.

If $c = (\mathbb{H} \setminus K, x, y = \infty, \dots)$, $h_*c = (\mathbb{H} \setminus K', x', y' = \infty, \dots)$ are equivalent configurations, $h'(\infty) = 1$, and we define $b = \kappa \partial_x \psi / \psi$, we get:

$$b(h_*c) = \frac{\kappa}{h'(x)} \frac{\partial_x(h'(x)^{-\alpha}\psi(c))}{h'(x)^{-\alpha}\psi(c)} = \frac{b(c)}{h'(x)} - \kappa \alpha \frac{h''}{h'^2}(x)$$

where c is implicitly a function of x (everything else being fixed). So if $\alpha = \alpha_\kappa = h_{1;2}(\kappa) = (6 - \kappa)/2\kappa$, one can define an SLE starting from the α -covariant partition function ψ (at least up to some positive stopping time). We denote this by $\text{SLE}_\kappa(\psi)$. This is well-defined for some positive time; we will not consider here the (difficult) questions of long-time behaviour.

9.1 Commutation conditions

Now assume we are given two α_κ -covariant functions ψ_1, ψ_2 on the configuration space, and use them to define SLEs starting resp. at x and y . purposes). In a configuration $c = (\mathbb{H} \setminus K, x, y, \dots)$, one grows SLEs at x and y up to capacity $\varepsilon, \varepsilon'$ (seen from infinity in $\mathbb{H}$; this is also arbitrary) and consider the effect on functions of x and y (after erasing hulls using chordal SLE). As earlier, this leads to the commutation conditions on ψ_1, ψ_2 . To make the argument neater, we compute on a section of the moduli space, as in [8, 2]. As before,

there is a marked point on the boundary used for normalization. In fact, in this chordal setup, it is more convenient not to quotient by automorphisms that fix this point; so all the construction will commute with scaling and translation. For definiteness, consider the following type of configurations: the upper half-plane $\mathbb{H}$ minus horizontal slits with appropriate marked points (including ∞). The only equivalences between such configurations are given by scaling and translation. Let $\mathfrak{D}$ be this family.

Consider an element $D_0 = (H_0, x, y, \dots) \in \mathfrak{D}$, that is H_0 is the half-plane minus some horizontal slits. We grow an $\text{SLE}_\kappa(\psi_1)$ at x up to half-plane capacity ε , then an $\text{SLE}_\kappa(\psi_2)$ at y (in the remaining domain) stopped when it reaches capacity ε (in the original domain). We then revert the order of the procedure and look for necessary conditions for these two procedures to yield the same distribution (of configuration). In particular, we compare the effect on moduli, up to second order in ε .

So consider a Loewner chain (K_t) growing at x . Let g_t be a conformal equivalence $\mathbb{H} \setminus K_t \rightarrow \mathbb{H}$, and f_t a conformal equivalence $D_0 \setminus K_t \rightarrow D_t$ for some $D_t \in \mathfrak{D}$. Everything is uniquely defined if we impose hydrodynamic normalization at infinity for g_t, f_t . Then let $h_t = f_t \circ g_t^{-1}$. By construction, g_t solves the Loewner equations:

$$\partial_t g_t = \frac{2}{g_t - X_t}, dX_t = \sqrt{\kappa} dB_t + \kappa \frac{\partial_x \psi_1(H_t)}{\psi_1(H_t)} dt$$

where ψ_1 is evaluated at the configuration $H_t = (g_t(H_0 \setminus K_t), X_t, \dots)$. Now consider $(\partial_t f_t) \circ f_t^{-1}$. This is a meromorphic function on D_t , taking real values on $\mathbb{R}$, vanishing at ∞ , regular except at x where it has a simple pole, and with constant imaginary part on the slits. It is easy to see that for each $D \in \mathfrak{D}$, there is a unique function V_D (Schwarz kernel) satisfying these conditions and with residue 2 at x . Also define A_D, B_D by:

$$V_D(w) = \frac{2}{w - x} + A_D + B_D(w - x) + O((w - x)^2)$$

Then:

$$\begin{aligned} \partial_t f_t &= h'_t(X_t)^2 V_{D_t} \circ f_t \\ \partial_t h_t(w) &= \partial_t (f_t \circ g_t^{-1})(w) = h'_t(X_t)^2 V_{D_t} \circ h_t(w) - \frac{2h'_t(w)}{w - X_t} \\ \partial_t h'_t(w) &= h'_t(X_t)^2 \cdot (h'_t \cdot V'_{D_t} \circ h_t)(w) - \frac{2h''_t(w)}{w - X_t} + \frac{2h'_t(w)}{(w - X_t)^2}. \end{aligned}$$

Taking limits at $w = X_t$, one gets:

$$\begin{aligned} (\partial_t h_t)(X_t) &= h'_t(X_t)^2 A_{D_t} - 3h''_t(X_t) \\ (\partial_t h'_t)(X_t) &= h'_t(X_t)^3 B_{D_t} + \left(\frac{h''_t}{2h'_t} - \frac{4h'''_t}{3} \right) (X_t) \end{aligned}$$

which gives in particular

$$\begin{aligned} dh_t(X_t) &= h'_t(X_t) dX_t + \left(\frac{\kappa}{2} - 3 \right) h''_t(X_t) dt + h'_t(X_t)^2 A_{D_t} dt \\ dh'_t(X_t) &= h''_t(X_t) dX_t + \left(\frac{h''_t(X_t)^2}{2h'_t(X_t)} + \left(\frac{\kappa}{2} - \frac{4}{3} \right) h'''_t(X_t) \right) dt + h'_t(X_t)^3 B_{D_t} dt \end{aligned}$$

as in [13] (and [8] for the case of annuli, using the explicit Villat kernel). Also:

$$\partial_t f'_t(y) = h'_t(X_t)^2 \cdot (f'_t \cdot V'_{D_t} \circ f_t)(y).$$

Now $\mathfrak{D}$ can be parametrized by a list of complex numbers $(x, y, z_1, \dots, z_m)$ containing marked points on $\mathbb{R}$, other marked points (including endpoints of horizontal slits) and their conjugates. Then the infinitesimal generator at D is:

$$\mathcal{L}_1 = \frac{\kappa}{2} \partial_{xx} + \left(A_D + \kappa \frac{\partial_x \psi_1}{\psi_1} \right) \partial_x + V_D(y) \partial_y + \sum_i V_D(z_i) \partial_{z_i}$$

with $V_D = V_{(x,y,z_1,\dots)}$ and ψ_1 is evaluated at D ; the restriction of ψ_1 to $\mathfrak{D}$ can also be seen as a function of $(x, y, z_1, \dots)$. The generator $\mathcal{L}_2$ associated with the $\text{SLE}_\kappa(\psi_2)$ growing at y is derived in similar fashion, and we get the commutation condition:

$$[\mathcal{L}_1, \mathcal{L}_2] = 2V'_y(x)\mathcal{L}_1 - 2V'_x(y)\mathcal{L}_2$$

where $V_x(z) = 2/(z-x) + \dots$ and $V_y(z) = 2/(z-y) + \dots$ are the Schwarz kernels with poles at x and y respectively (and depend implicitly on the other moduli). We will use the notation:

$$\ell_x = V_x(y)\partial_y + \sum V_x(z_i)\partial_i, \quad \ell_y = V_y(x)\partial_x + \sum V_y(z_i)\partial_i$$

for the part that does not involve κ, ψ_1, ψ_2 .

Expanding the commutation condition, we get the equations:

$$\partial_{xy} \log(\psi_1/\psi_2) = 0 \tag{9.4}$$

and:

$$\begin{aligned} -\mathcal{L}_2 \left(A_x + \kappa \frac{\partial_x \psi_1}{\psi_1} \right) + \mathcal{L}_1(V_y(x)) + 2V'_x(y)V_y(x) - 2V'_y(x) \left(A_x + \kappa \frac{\partial_x \psi_1}{\psi_1} \right) &= 0 \\ -\mathcal{L}_1 \left(A_y + \kappa \frac{\partial_y \psi_2}{\psi_2} \right) + \mathcal{L}_2(V_x(y)) + 2V'_y(x)V_x(y) - 2V'_x(y) \left(A_y + \kappa \frac{\partial_y \psi_2}{\psi_2} \right) &= 0 \end{aligned} \tag{9.5}$$

All other conditions are identities not involving ψ_1, ψ_2 :

$$(\ell_x + 2V'_x(y))V_y(z) = (\ell_y + 2V'_y(x))V_x(z) \tag{9.6}$$

For this, note that $V_y(z)$ does not depend on x and other marked points, but depends on endpoints of horizontal slits. Considering the difference between left-hand side and right-hand side as a function of z , in particular its expansion at x, y , one sees that it extends to a bounded holomorphic function on the Schottky double of D and vanishes at infinity, hence is identically 0. The probabilistic interpretation of the Schwarz kernel V given in [14] can also be used to prove this identity, along the lines of Lemma 14 (ii) below.

Consider the conditions (9.4),(9.5). From (9.4), one can write $\psi_1 = \psi_2 = \psi$, by multiplying ψ_1 by a function that does not depend on x (so that the definition of $\text{SLE}_\kappa(\psi_1)$ is not affected), and doing the same for ψ_2 . One can write:

$$\begin{aligned} \mathcal{L}_1 \left(\frac{\partial_y \psi}{\psi} \right) - \partial_y \left(\frac{\mathcal{L}_1 \psi}{\psi} \right) &= \frac{[\mathcal{L}_1, \partial_y] \psi}{\psi} + \kappa \frac{\partial_y \psi (\partial_x \psi)^2}{\psi^3} - \kappa \frac{\partial_x \psi \partial_{xy} \psi}{\psi^2} \\ &= -V'_x(y) \partial_y \log(\psi) - \kappa \partial_{xy} \log(\psi) \partial_x \log(\psi) + \kappa \partial_y \log(\psi) (\partial_x \log(\psi))^2 - \kappa \frac{\partial_x \psi \partial_{xy} \psi}{\psi^2} \\ &= -V'_x(y) \partial_y \log(\psi) - \kappa \partial_y (\partial_x \log \psi)^2 \end{aligned}$$

Let $\check{\mathcal{L}}_1$ be the differential operator obtained by setting $\psi_1 = 1$ in $\mathcal{L}_1$:

$$\check{\mathcal{L}}_1 = \frac{\kappa}{2} \partial_{xx} + A_x \partial_x + V_D(y) \partial_y + \sum_i V_D(z_i) \partial_{z_i} = \frac{\kappa}{2} \partial_{xx} + A_x \partial_x + \ell_x$$

and $\check{\mathcal{L}}_2$ is defined in the same fashion. Then (9.5) can be written as:

$$\kappa \partial_y \left(\frac{\check{\mathcal{L}}_1 \psi}{\psi} \right) = \check{\mathcal{L}}_2(V_x(y)) - \check{\mathcal{L}}_1(A_y) + 2V'_y(x)V_x(y) - 2A_y V'_x(y)$$

and similarly for the other condition. Consider now (9.6), applied to a marked point $z = y + \varepsilon$:

$$\begin{aligned}
0 &= V_x(y)\partial_y V_y(z) + V_x(z)\partial_z V_y(z) + \sum_i V_x(z_i)\partial_{z_i} V_y(z) - V_y(x)\partial_x V_x(z) - V_y(z)\partial_z V_x(z) - \sum_i V_y(z_i)\partial_{z_i} V_x(z) \\
&\quad + 2V'_x(y)V_y(z) - 2V'_y(x)V_x(z) \\
&= V_x(y)\left(\frac{2}{\varepsilon^2} + \partial_y A_y - B_y + (\partial_y B_y - 2C_y)\varepsilon\right) + \left(V_x(y) + \varepsilon V'_x(y) + \frac{\varepsilon^2}{2}V''_x(y) + \frac{\varepsilon^3}{6}V'''_x(y)\right)\left(-\frac{2}{\varepsilon^2} + B_y + 2C_y\varepsilon\right) \\
&\quad + \sum_i V_x(z_i)\partial_{z_i}(A_y + \varepsilon B_y) - V_y(x)\partial_x V_x(y) - V_y(x)\partial_x V'_x(y)\varepsilon - \left(\frac{2}{\varepsilon} + A_y + B_y\varepsilon\right)\left(V'_x(y) + \varepsilon V''_x(y) + \frac{\varepsilon^2}{2}V'''_x(y)\right) \\
&\quad - \sum_i V_y(z_i)\partial_{z_i}(V_x(y) + \varepsilon V'_x(y)) + 2V'_x(y)\left(\frac{2}{\varepsilon} + A_y + B_y\varepsilon\right) - 2V'_y(x)V_x(y) - 2V'_y(x)V'_x(y)\varepsilon + O(\varepsilon^2)
\end{aligned}$$

where C_y is such that $V_y(z) = 2/\varepsilon + A_y + B_y\varepsilon + C_y\varepsilon^2 + O(\varepsilon^3)$. Considering the coefficients of ε^0 , ε^1 , it follows that:

$$(\ell_x + 2V'_x(y))(A_y) - (\check{\mathcal{L}}_2 + 2V'_y(x))(V_x(y)) + \left(\frac{\kappa}{2} - 3\right)V''_x(y) = 0 \quad (9.7)$$

and

$$(\ell_x + 2V'_x(y))(B_y) - (\check{\mathcal{L}}_2 + 2V'_y(x))(V'_x(y)) + \left(\frac{\kappa}{2} - \frac{4}{3}\right)V'''_x(y) = 0 \quad (9.8)$$

Of course one can exchange the roles of x and y in these identities. So (9.5) can be written as:

$$\kappa\partial_y\left(\frac{\check{\mathcal{L}}_1\psi}{\psi}\right) = \left(\frac{\kappa}{2} - 3\right)V''_x(y)$$

or (with the symmetric condition):

$$\partial_y\left(\frac{(\check{\mathcal{L}}_1 + \alpha_\kappa V'_x(y))\psi}{\psi}\right) = \partial_x\left(\frac{(\check{\mathcal{L}}_2 + \alpha_\kappa V'_y(x))\psi}{\psi}\right) = 0.$$

9.2 Restriction martingales

Restriction-like martingales in multiply-connected domains involving harmonic invariants are studied in [14, 27], in particular when $\kappa = 2$. When $\kappa \neq 2$, these “harmonic” martingales are distinct from those we are discussing in these sections, though methods are fairly similar.

Let D be a subdomain of $\mathbb{H}$ (which we can freely assume if we want to define Möbius invariant distributions); the boundary of D contains open real segments around $x, y \in \mathbb{R}$. Define: $\Gamma(D, x) = \mu_x^{bub}(\{\delta \subsetneq D\})$, where μ_x^{bub} is the bubble measure rooted at x (see [17]). Then:

$$\Gamma(D, x) = \partial_{n_y}(M_{\mathbb{H}}(y, x) - M_D(y, x))|_{y=x}$$

where M_D denotes the minimal function with singularity at x , i.e. the positive harmonic function in D that extends continuously to 0 on the boundary except at x , with normalization $M_D(x + i\varepsilon, x) = \varepsilon^{-1}(1 + O(\varepsilon))$. This function can be obtained by taking the normal derivative at x of the Green’s function $G_D(y, x)$ (with adequate normalization). Let ϕ be a conformal equivalence $D \rightarrow D' = \phi(D)$. From the conformal invariance property of the Green’s function, it is easy to derive the covariance property of the minimal function:

$$M_D(y, x) = \phi'(x)M_{\phi(D)}(\phi(y), \phi(x))$$

and then the Schwarzian-like covariance property for Γ :

$$\Gamma(D, x) = \phi'(x)^2\Gamma(\phi(D), \phi(x)) - \frac{S\phi(x)}{6}.$$

Note that in the case where D is simply connected, $\Gamma(D, x) = -S\phi_D(x)/6$, and this is the usual covariance property of Schwarzian derivatives:

$$S(\phi_D) = S(\phi_{D'} \circ \phi) = (\phi')^2 (S\phi_{D'}) \circ \phi + S\phi$$

We are interested in the following situation: let γ be a chordal SLE_κ in $\mathbb{H}$, say $\kappa \leq 8/3$, and L is an independent loop soup with intensity λ_κ . Condition on the event that $\gamma^L = \gamma \cup \{\delta \in L : \delta \cap \gamma \neq \emptyset\}$ stays in D_0 . It is not clear at this point that the probability of this event is α_κ -covariant, hence that the resulting distribution is conformally invariant.

First consider a chordal SLE_κ from x to y in $\mathbb{H} \supset D_0$ (a chordal SLE unaware of the presence of holes), and an α_κ -covariant function φ on the moduli space. By Möbius invariance, one can send y to infinity (and use hydrodynamic normalization); so $(X_t/\sqrt{\kappa})$ is now a standard Brownian motion. Here $g_t : D_0 \setminus K_t \rightarrow H_t$ is a conformal equivalence that extends through the holes, $h_t : H_t \rightarrow D_t$ is a conformal equivalence and $D_t \in \mathfrak{D}$ is a standard domain. Let $M_t = \varphi(D_0 \setminus K_t, \dots)$. From the covariance assumption,

$$M_t = h'_t(X_t)^\alpha \varphi(h_t(X_t), \dots).$$

Set $\alpha = \alpha_\kappa = (6 - \kappa)/2\kappa$. Then:

$$\frac{dh'_t(X_t)^\alpha}{h'_t(X_t)^\alpha} = \alpha \frac{h''_t(X_t)}{h'_t(X_t)} dX_t - \frac{\lambda_\kappa}{6} Sh_t(X_t) dt + \alpha h'_t(X_t)^2 B_{D_t} dt$$

where $\lambda_\kappa = (8 - 3\kappa)(6 - \kappa)/2\kappa$ and Sh_t is the Schwarzian derivative. Besides:

$$\begin{aligned} d\varphi(h_t(X_t), \dots, f_t(z_i), \dots) = & h'_t(X_t) \partial_x \varphi dX_t + \left[\frac{\kappa}{2} h'_t(X_t)^2 \partial_{x,x} + \left(\frac{\kappa}{2} - 3 \right) h''_t(X_t) \partial_x \right. \\ & \left. + h'_t(X_t)^2 A_{D_t} \partial_x + h'_t(X_t)^2 \sum_i V_{D_t}(z_i) \partial_{z_i} \right] \varphi(h_t(X_t), \dots, f_t(z_i), \dots) dt \end{aligned}$$

Let:

$$Z_t = M_t \exp \left(-\lambda_\kappa \int_0^t \Gamma(H_s, X_s) ds \right) = M_t \exp \left(-\lambda_\kappa \int_0^t \left(h'_s(X_s)^2 \Gamma(D_s, h_s(X_s)) - \frac{Sh_s(X_s)}{6} \right) ds \right)$$

Then Z_t is a local martingale iff φ (restricted to $\mathfrak{D}$, hence considered as a function of the parameters $x, \dots, z_i, \dots$) is annihilated by the differential operator:

$$\mathcal{M} = \frac{\kappa}{2} \partial_{xx} + A_D \partial_x + \ell_x + \alpha B_D - \lambda_\kappa \Gamma_x$$

where $\Gamma_x = \Gamma(D, x)$ in the standard domain D . If y is finite (and another marked point is used for normalization), one can compute along the same lines. So if C_t is the covariance factor:

$$C_t = \left(h'_t(X_t) h'_t(Y_t) \left(\frac{Y_t - X_t}{h_t(Y_t) - h_t(X_t)} \right)^2 \right)^\alpha$$

then

$$\begin{aligned}
\frac{dC_t}{C_t} &= \left(\alpha \frac{h_t''(X_t)}{h_t'(X_t)} dX_t - \frac{\lambda_\kappa}{6} Sh_t(X_t) dt + \alpha h_t'(X_t)^2 B_{D_t} dt \right) \\
&\quad + \alpha \left(h_t'(X_t)^2 V_{D_t}'(h_t(Y_t)) + \frac{2}{(Y_t - X_t)^2} \right) dt + 2\alpha \left(\frac{dX_t}{X_t - Y_t} + (5 - \kappa) \frac{dt}{(Y_t - X_t)^2} \right) \\
&\quad - \frac{2\alpha}{h_t(X_t) - h_t(Y_t)} \left(h_t'(X_t) dX_t + \left(\frac{\kappa}{2} - 3 \right) h_t''(X_t) dt + h_t'(X_t)^2 (A_{D_t} - V_{D_t}(h_t(Y_t))) - \frac{3}{h_t(X_t) - h_t(Y_t)} dt \right) \\
&\quad + 2\kappa\alpha^2 \left(\frac{h_t''(X_t)}{h_t'(X_t)(X_t - Y_t)} - \frac{h_t''(X_t)}{h_t(X_t) - h_t(Y_t)} - 2 \frac{h_t'(X_t)}{(X_t - Y_t)(h_t(X_t) - h_t(Y_t))} \right) dt \\
&= \alpha \left(\frac{h_t''(X_t)}{h_t'(X_t)} + \frac{2}{X_t - Y_t} - \frac{2h_t'(X_t)}{h_t(X_t) - h_t(Y_t)} \right) \sqrt{\kappa} dB_t - \frac{\lambda_\kappa}{6} Sh_t(X_t) dt \\
&\quad + \alpha h_t'(X_t)^2 \left(B_{D_t} + V_{D_t}'(h_t(Y_t)) - \frac{2}{h_t(X_t) - h_t(Y_t)} (A_{D_t} - V_{D_t}(h_t(Y_t))) - \frac{3}{h_t(X_t) - h_t(Y_t)} \right) dt
\end{aligned}$$

So M_t is a local martingale iff φ is annihilated by the operator:

$$\mathcal{M} = \frac{\kappa}{2} \partial_{xx} + (A_D + \frac{\kappa - 6}{x - y}) \partial_x + \ell_x + \alpha \left(B_D + V_D'(y) + \frac{2(V_D(y) - A_D)}{x - y} + \frac{6}{(x - y)^2} \right) - \lambda_\kappa \Gamma_x$$

Observe that:

$$\begin{aligned}
(x - y)^{-2\alpha} \mathcal{M}(x - y)^{2\alpha} &= \frac{\kappa}{2} \partial_{xx} + A_D \partial_x + \ell_x + \alpha (B_D + V_D'(y)) + \lambda_\kappa \Gamma_x \\
&= \tilde{\mathcal{L}}_1 + \alpha (B_D + V_D'(y)) - \lambda_\kappa \Gamma_x
\end{aligned}$$

The conjugation corresponds to a change of reference measure (SLE aiming at ∞ rather than SLE aiming at y).

We sum up the discussion of this section. As before, $\mathfrak{C}$ is a configuration space, x and y are marked points on a boundary component of a configuration; $\mathfrak{M}$ is the associated moduli space, and $\mathfrak{D}$ is a class of standard domains, with associated Schwarz kernels V . We note V_x, V_y to emphasize the pole of the kernel.

Proposition 13. (i) Let ψ_1, ψ_2 be α_κ -covariant functions; consider an $\text{SLE}_\kappa(\psi_1)$ growing at x and an $\text{SLE}_\kappa(\psi_2)$ growing at y . These two SLEs satisfy commutation condition iff there is an α_κ -covariant function ψ such that $\text{SLE}_\kappa(\psi_i)$, $i = 1, 2$, is distributed as an $\text{SLE}_\kappa(\psi)$ growing at x (resp. y) and ψ satisfy the conditions:

$$\partial_y \left(\frac{(\tilde{\mathcal{L}}_1 + \alpha V_x'(y)) \psi}{\psi} \right) = \partial_x \left(\frac{(\tilde{\mathcal{L}}_2 + \alpha V_y'(x)) \psi}{\psi} \right) = 0$$

(ii) Let φ be an α_κ -covariant function. Let $c \in \mathfrak{C}$ be a configuration. One can assume that $c = D_0$ is a standard domain. Consider a chordal SLE_κ from x to y in c with holes erased; let $(c_t)_t$ be the corresponding family of configurations. Let h_t be the map from c_t to a standard domain $D_t \in \mathfrak{D}$, with hydrodynamic normalization. Let:

$$Z_t = \varphi(c_t) \exp \left(-\lambda_\kappa \int_0^t \Gamma(c_s, \gamma_s) ds \right)$$

Then Z is a local martingale iff φ restricted to $\mathfrak{D}$ is annihilated by the operator:

$$\mathcal{M} = (x - y)^{2\alpha} (\tilde{\mathcal{L}}_1 + \alpha (B_x + V_x'(y)) - \lambda_\kappa \Gamma_x) (x - y)^{-2\alpha}$$

(iii) In the situation of (ii), consider the Girsanov transform of chordal SLE by Z . Then the resulting process is an $\text{SLE}_\kappa(\psi)$, where ψ is an α_κ -covariant function whose restriction to $\mathfrak{D}$ is annihilated by the operator:

$$\tilde{\mathcal{L}}_1 + \alpha_\kappa (B_x + V_x'(y)) - \lambda_\kappa \Gamma_x$$

In particular, $\text{SLE}_\kappa(\psi)$ started at x and $\text{SLE}_\kappa(\psi)$ started at y satisfy the commutation conditions.

9.3 The case $\kappa = 0$

In this section, we address the following (non trivial) question: how to define a (deterministic) reversible SLE_0 in a multiply connected domain ? This requires implicitly the domain Markov property and conformal invariance. While there are many ways to construct conformally invariant chords (e.g. as level lines or flow lines of harmonic invariants), it is not so obvious to satisfy all conditions simultaneously. We propose here a construction based on the restriction ideas (and do not claim to prove anything rigorous about it).

Consider a nice subdomain H_0 of $\mathbb{H}$ (i.e. $\overline{\mathbb{H} \setminus H_0}$ is compact and has finitely many components), $x_0 \in \mathbb{R}$, $(x_0 - \varepsilon, x_0 + \varepsilon) \subset \partial H_0$ for some $\varepsilon > 0$; we attempt to describe an $\text{SLE}_0(x_0 \rightarrow \infty)$ in H_0 . The idea is to take γ a chordal $\text{SLE}_\kappa(x_0 \rightarrow \infty)$ in $\mathbb{H}$ conditionally on γ^L staying in H_0 , where L is an independent loop soup with intensity λ_κ , and take the limit as $\kappa \searrow 0$. On the one hand, the unconditional chordal SLE_κ converges to a hyperbolic geodesic in $\mathbb{H}$, which is not conformally invariant; on the other hand the intensity of the loop soup diverges. We shall define a total cost summing the large deviation rate for Brownian motion and the loop soup term, which will give a conformally covariant functional.

To check conformal invariance, we will use comparisons of Loewner chains in conformally equivalent domains as in [13] (see also the previous subsections). Let us fix some notations. Let $\phi_0 : H_0 \rightarrow \tilde{H}_0$ be a conformal equivalence to $\tilde{H}_0$ (and $\tilde{H}_0$ satisfies the same properties as H_0). Consider the Loewner flow:

$$\partial_t g_t = \frac{2}{g_t - X_t}$$

that maps H_0 to H_t (and extends to $\mathbb{H}$). This Loewner chain is mapped by ϕ_0 to a Loewner chain corresponding to the flow:

$$\partial_t \tilde{g}_t = \frac{2\phi'_t(X_t)^2}{\tilde{g}_t - \tilde{X}_t}$$

where $\phi_t = \tilde{g}_t \circ \phi_0 \circ g_t^{-1}$ is a conformal equivalence $H_t \rightarrow \tilde{H}_t$ (and $\tilde{X}_t = \phi_t(X_t)$ etc...).

Consider now the functional on driving processes:

$$I(X) = I(X, H_0) = \frac{1}{2} \int_0^\infty (\partial_t X_t)^2 dt + 24 \int_0^\infty \Gamma(H_t, X_t) dt.$$

Informally, as $\kappa \searrow 0$, a driving process X has weight $\propto \exp(-I(X)/\kappa)$, the first term being the large deviations rate for Brownian motion, the second term coming from the loop soup conditioning ($\lambda_\kappa \sim 24/\kappa$). We want to prove that $I(X) = I(\tilde{X}) + c$, where $\tilde{X}$ has time parameter:

$$s(t) = \int_0^t \phi'_u(X_u)^2 du.$$

Taking into account the time change, we get:

$$I(\tilde{X}) = I(\tilde{X}, \tilde{H}_0) = \frac{1}{2} \int_0^\infty \left(\frac{\partial_t \tilde{X}_t}{\phi'_t(X_t)^2} \right)^2 \phi'_t(X_t)^2 dt + 24 \int_0^\infty \phi'_t(X_t)^2 \Gamma(\tilde{H}_t, \tilde{X}_t) dt.$$

From the covariance of Γ , we have:

$$\Gamma(H_t, X_t) = \phi'_t(X_t)^2 \Gamma(\tilde{H}_t, \tilde{X}_t) - \frac{(S\phi_t)(X_t)}{6}.$$

Also, comparisons of Loewner chains yield:

$$\begin{aligned} \partial_t \tilde{X}_t &= \phi'_t(X_t) \partial_t X_t - 3\phi''_t(X_t) \\ \partial_t(\phi'_t(X_t)) &= \phi''_t(X_t) \partial_t X_t + \left(\frac{\phi''_t}{2\phi'_t} - \frac{4}{3}\phi'''_t \right) (X_t) \end{aligned}$$

It follows that:

$$\begin{aligned}
I(\tilde{X}) - I(X) &= \frac{1}{2} \int_0^\infty \left(-6 \frac{\phi_t''}{\phi_t'}(X_t) \partial_t X_t + 9 \left(\frac{\phi_t''}{\phi_t'} \right)^2 (X_t) \right) dt + 4 \int_0^\infty S \phi_t(X_t) dt \\
&= \frac{1}{2} \int_0^\infty \left(-6 \frac{\phi_t''}{\phi_t'}(X_t) \partial_t X_t - 3 \left(\frac{\phi_t''}{\phi_t'} \right)^2 (X_t) + 8 \frac{\phi_t'''}{\phi_t'}(X_t) \right) dt = -3 \int_0^\infty \frac{\partial_t(\phi_t'(X_t))}{\phi_t'(X_t)} dt \\
&= 3 \log(\phi_0'(x_0))
\end{aligned}$$

since $\phi_t'(X_t) \rightarrow 1$ (ϕ_t has hydrodynamic normalization at ∞). Thus we get the covariance relation:

$$I(\phi_* X, \phi_* H) = I(X, H) + 3 \log(\phi'(x_0)).$$

So provided that I attains a minimum, the minimizing path is conformally invariant and Markov. Also, one can minimize over a subset of paths satisfying particular global topological conditions (e.g. leaving a hole on the left). Assuming that chordal SLE_κ in $\mathbb{H}$ is reversible, this is also reversible (the conditioning event does not depend on the orientation of the paths). For a configuration (H, x, ∞) , denote:

$$\psi(H) = \sup_X \left(\exp^{-I(H, X)} \right)$$

which is (-3) -covariant. With the previous notations, one can write the coherence condition:

$$\psi(H) = \sup_{X_{[0,t]}} \left(\psi(H_t) \exp \left(-\frac{1}{2} \int_0^t (\partial_s X_s)^2 ds + 24 \int_0^t \Gamma(H_s, X_s) ds \right) \right)$$

By conformal invariance, one can assume that H is a standard configuration. Let $h_t : H_t \rightarrow D_t$ be the normalized conformal equivalence to a standard domain. Then:

$$\psi(D_t) = \psi(H_t) (h_t'(X_t))^{-3}$$

and $\partial_t(h_t(X_t))|_{t=0} = A_x$, $\partial_t(h_t'(X_t))|_{t=0} = B_x$. Let $\overline{X}$ denote a maximizing driving function. Expanding the previous condition for small t , one gets:

$$(\ell_x + A_x \partial_x + (\partial_t \overline{X}_t) \partial_x + 3B_x) \psi - \left(\frac{1}{2} (\partial_t \overline{X}_t)^2 + 24 \Gamma_x \right) \psi = 0$$

at time $t = 0$ (where ψ is seen as a function on standard configurations). Besides, the left-hand side is optimal for $\partial_t \overline{X}_t$. This implies that:

$$\partial_t \overline{X}_t = \frac{\partial_x \psi}{\psi} \quad \text{and} \quad \frac{(\partial_x \psi)^2}{2\psi} + (\ell_x + A_x \partial_x + 3B_x - 24 \Gamma_x) \psi = 0.$$

One can also get this equation by considering:

$$\left(\frac{\kappa}{2} \partial_{xx} + A_x \partial_x + \ell_x + \alpha_\kappa B_x - \lambda_\kappa \Gamma_x \right) \psi_\kappa = 0$$

and then rewrite the equation for $(\psi_\kappa)^\kappa$ and take the limiting equation as $\kappa \searrow 0$. Now if the target point is not ∞ but a finite point y , one can proceed similarly. The condition obtained on ψ for growth at x, y gives the commutation condition.

9.4 Towards a classification

In simply connected domains, we obtained a complete classification of commuting SLEs. In the multiply connected case, it appears to be much more technical, so we shall only outline some elements. First, if we don't assume *a priori* that the two SLEs have the same κ , then it is not hard to check that the commutation condition for an $\text{SLE}_\kappa(\psi)$ growing at x and an $\text{SLE}_{\tilde{\kappa}}(\psi)$ growing at y writes:

$$\partial_y \left(\frac{(\check{\mathcal{L}}_1 + \alpha_{\tilde{\kappa}} V'_x(y)) \psi}{\psi} \right) = \partial_x \left(\frac{(\check{\mathcal{L}}_2 + \alpha_\kappa V'_y(x)) \psi}{\psi} \right) = 0$$

where now $\check{\mathcal{L}}_2 = \frac{\tilde{\kappa}}{2} \partial_{yy} + \dots$. Also, if $\tilde{\kappa} \neq \kappa$, the covariance condition for ψ is modified as follows:

- (i). (Möbius invariance) For any $c = (D \setminus K, \dots)$, $c' = (D' \setminus K', \dots)$ where D, D' are simply connected, $h : c \rightarrow c' = h_* c$ is an equivalence of configurations, and h extends to a conformal equivalence $D \rightarrow D'$, one has:

$$\psi(h_* c) = \psi(c)$$

- (ii). (covariance) For any $c = (\mathbb{H} \setminus K, x, x', y, y', \dots)$, $c' = (\mathbb{H} \setminus K', \dots)$ where D is simply connected, $x, y \in \partial D$, ∂D is smooth at x, y , and $h : c \rightarrow c' = h_* c$ is an equivalence of configurations, one has:

$$\psi(h_* c) = \left(h'(x) h'(x') \left(\frac{x - x'}{h(x) - h(x')} \right)^2 \right)^{-\alpha_\kappa} \left(h'(y) h'(y') \left(\frac{y - y'}{h(y) - h(y')} \right)^2 \right)^{-\alpha_{\tilde{\kappa}}} \psi(c)$$

where x', y' are new target points for the SLEs.

We now revert to the case $\kappa = \tilde{\kappa}$ (and the two SLEs are “aiming at each other”) as discussed earlier and further study the commutation conditions. Consider the operators:

$$\mathcal{M}_1 = \check{\mathcal{L}}_1 + \alpha V'_x(y) + h_1, \mathcal{M}_2 = \check{\mathcal{L}}_2 + \alpha V'_y(x) + h_2$$

where $\partial_y h_1 = \partial_x h_2 = 0$. To define two commuting SLEs, we have to find functions h_1, h_2, ψ such that $\mathcal{M}_1 \psi = \mathcal{M}_2 \psi = 0$.

Say $\kappa = 8/3$, and consider a chordal $\text{SLE}_{8/3}$ in a simply connected domain conditioned to avoid some holes. Then the conditional SLE can be represented as an $\text{SLE}_{8/3}(\psi)$, and ψ (restricted to a section of the moduli space) is annihilated by a differential operator, coming from the restriction property. Since $\text{SLE}_{8/3}$ is revertible, so is the conditional version; this gives a commutation condition in a multiply connected domain. For SLE_2 , we know that the restriction construction is the scaling limit of a revertible discrete model, hence the restriction weight $h_1(x) = \alpha_2 B_x - \lambda_2 \Gamma_x$ should be a solution of this functional equation. We now give a direct derivation of this fact.

Lemma 14. (i). *If there exists a non-vanishing function ψ such that $\mathcal{M}_1 \psi = \mathcal{M}_2 \psi = 0$, then h_1, h_2 satisfy:*

$$(\check{\mathcal{L}}_1 + 2V'_x(y))(\alpha V'_y(x) + h_2) = (\check{\mathcal{L}}_2 + 2V'_y(x))(\alpha V'_x(y) + h_1)$$

(ii). *The following identity holds:*

$$(\ell_x + 2V'_x(y))\Gamma_y - \frac{V_x'''(y)}{6} = (\ell_y + 2V'_y(x))\Gamma_x - \frac{V_y'''(x)}{6}.$$

(iii). *The condition (i) is satisfied if $h_1(x, \dots) = h(x, \dots)$, $h_2(y, \dots) = h(y, \dots)$, and*

$$h(x, \dots) = \alpha_\kappa B_x - \lambda_\kappa \Gamma_x + \ell_x f$$

where f is a function on standard configurations that does not depend on x, y .

Proof. (i) If ψ is such that $\mathcal{M}_1\psi = \mathcal{M}_2\psi = 0$, then also $\mathcal{M}_0\psi = 0$ where:

$$\begin{aligned}\mathcal{M}_0 &= [\mathcal{M}_1, \mathcal{M}_2] - 2V'_y(x)\mathcal{M}_1 + 2V'_x(y)\mathcal{M}_2 \\ &= (\check{\mathcal{L}}_1 V_y(x) - \check{\mathcal{L}}_2 A_x + 2V'_x(y)V_y(x) - 2A_x V'_y(x) + \kappa \alpha V''_y(x)) \partial_x \\ &\quad - (\check{\mathcal{L}}_2 V_x(y) - \check{\mathcal{L}}_1 A_y + 2V'_y(x)V_x(y) - 2A_y V'_x(y) + \kappa \alpha V''_x(y)) \partial_y \\ &\quad + \sum (\check{\mathcal{L}}_1 V_y(z_i) - \check{\mathcal{L}}_2 V_x(z_i) + 2V'_x(y)V_y(z_i) - 2V'_y(x)V_x(z_i)) \partial_{z_i} \\ &\quad + ((\check{\mathcal{L}}_1 + 2V'_x(y))(\alpha V'_y(x) + h_2) - (\check{\mathcal{L}}_2 + 2V'_y(x))(\alpha V'_x(y) + h_1))\end{aligned}$$

The first-order terms vanish, from (9.6), (9.7).

(ii) We want to interpret the left-hand side and the right-hand side of the equation in terms of Brownian measures. Let us start with the left-hand side. In the standard domain D , we grow a vertical slit γ at x . For small t , $H_t = D \setminus \gamma_{[0,t]}$ is conformally equivalent to a standard domain D_t ; the conformal equivalence f_t can be expanded in t as:

$$f_t(z) = z + tV_x(z) + o(t)$$

where V_x is the Schwarz kernel with pole at x in the domain D . Hence $f'_t(y)^2 = 1 + 2tV'_x(y) + o(t)$ and $Sf_t(y) = tV'''_x(y) + o(t)$. Let us consider $\Gamma(H_t, y)$. From the covariance property of Γ , we get:

$$\begin{aligned}\Gamma(H_t, y) - \Gamma(H_0, y) &= f'_t(y)^2 \Gamma(D_t, f_t(y)) - \frac{Sf_t(y)}{6} - \Gamma(H_0, y) \\ &= t \left((V_x(y) \partial_y + \sum_i V_x(z_i) \partial_{z_i} + 2V'_x(y)) \Gamma_y - \frac{V'''_x(y)}{6} \right) + o(t)\end{aligned}$$

Now $\Gamma(H_t, y) - \Gamma(H_0, y)$ is the measure for Brownian bubbles rooted at y of bubbles that intersect $\gamma_{[0,t]}$ as well as one of the holes. Consider $\tilde{\gamma}$ a vertical slit growing at y , and the loop soup measure:

$$\begin{aligned}\mu_{\mathbb{H}}^{loop}(\{\delta : \delta \cap \gamma \neq \emptyset, \delta \cap \tilde{\gamma} \neq \emptyset, \delta \subsetneq H_0\}) &= \mu_x^{bub}(\{\delta : \delta \cap \tilde{\gamma} \neq \emptyset, \delta \subsetneq H_0\})(t + o(t)) \\ &= \mu_y^{bub}(\{\delta : \delta \cap \gamma \neq \emptyset, \delta \subsetneq H_0\})(t + o(t))\end{aligned}$$

as follows from Proposition 11 in [17]. Here we are considering loops that intersect the two small slits $\gamma, \tilde{\gamma}$. We can root such loops either near x or y , which gives us the right-hand side and the left hand-side of the claimed identity.

(iii) As we did earlier, we can expand the condition (i) at $y = x$, writing $h_1 = \alpha B_x - \lambda_\kappa \Gamma_x + h(x, \dots)$, $h_2 = \alpha B_y - \lambda_\kappa \Gamma_y + h(y, \dots)$ for some unknown function h . The functional equation for h is the linear equation :

$$(\ell_x + 2V'_x(y))h(y) = (\ell_y + 2V'_y(x))h(x)$$

(from (9.8) and (ii)). Note that this equation does not involve κ . Also, using the identity 9.6, one can write:

$$(\ell_x + 2V'_x(y))\ell_y + (\dots)\partial_x = (\ell_y + 2V'_y(x))\ell_x + (\dots)\partial_y.$$

So if f does not depend on x, y , $h(x) = \ell_x f$ gives a solution of the functional equation. $\square$

Let us consider again the equation “without second member”:

$$(\ell_x + 2V'_x(y))h(y) = (\ell_y + 2V'_y(x))h(x)$$

appearing in (iii). From (9.6), we see that if z is a marked point, then $h(x) = V_x(z)$ is a solution for any marked point z . Also, differentiating the identity (9.6) w.r.t. z , two terms $V'_x(z)V'_y(z)$ cancel out, and we get:

$$(\ell_x + 2V'_x(y))V'_y(z) = (\ell_y + 2V'_y(x))V'_x(z)$$

So $h(x) = V'_x(z)$ is a solution. Similarly, $h(x) = (V_x(z_2) - V_x(z_1))/(z_2 - z_1)$ is also a solution. This corresponds to rational solutions in the simply connected case. It is not so clear what would be f in these cases.

As in Section 4, we can interpret a solution h as the tangent map of a cocycle. More precisely, if h is as above, then one can define a function C on hulls and residual configurations (configurations determined by the domain and the marked points $z_1, \dots$) such that:

- (i). For all hulls A, B , $C(A.B, z) = C(B, \phi_A(z))C(A, z)$.
- (ii). If A is a hull of half-plane capacity ε located at x , then $C(A, z) = 1 + 2\varepsilon h(x, z) + o(\varepsilon)$.

Some of the arguments we used in the simply connected case can be replicated here; though a general classification of such cocycles appears to be more difficult.

Acknowledgments. I wish to thank Greg Lawler and Wendelin Werner for stimulating and fruitful conversations.

References

- [1] M. Aizenman and A. Burchard. Hölder regularity and dimension bounds for random curves. *Duke Math. J.*, 99(3):419–453, 1999.
- [2] R. O. Bauer and R. M. Friedrich. On Chordal and Bilateral SLE in multiply connected domains.
- [3] V. Beffara. The dimension of SLE curves. *preprint, arXiv:math.PR/0211322*, 2002.
- [4] F. Camia and C. M. Newman. The Full Scaling Limit of Two-Dimensional Critical Percolation.
- [5] J. Cardy. Corrigendum: “Stochastic Loewner evolution and Dyson’s circular ensembles” [J. Phys. A **36** (2003), no. 24, L379–L386]. *J. Phys. A*, 36(49):12343, 2003.
- [6] J. Cardy. Stochastic Loewner evolution and Dyson’s circular ensembles. *J. Phys. A*, 36(24):L379–L386, 2003.
- [7] J. Dubédat. SLE(κ, ρ) martingales and duality. *Ann. Probab.*, to appear, 2003.
- [8] J. Dubédat. Critical percolation in annuli and SLE₆. *Comm. Math. Phys.*, 245(3):627–637, 2004.
- [9] J. Dubédat. Excursion decompositions for SLE and Watts’ crossing formula. *preprint, arXiv:math.PR/0405074*, 2004.
- [10] J. Dubédat. Euler integrals for commuting SLEs. *preprint, arXiv:math.PR/0507276*, 2005.
- [11] S. Fomin. Loop-erased walks and total positivity. *Trans. Amer. Math. Soc.*, 353(9):3563–3583 (electronic), 2001.
- [12] M. J. Kozdron and G. F. Lawler. Estimates of random walk exit probabilities and application to loop-erased random walk.
- [13] G. Lawler, O. Schramm, and W. Werner. Conformal restriction: the chordal case. *J. Amer. Math. Soc.*, 16(4):917–955 (electronic), 2003.
- [14] G. F. Lawler. The Laplacian- b random walk and the Schramm-Loewner evolution .

- [15] G. F. Lawler. *Conformally invariant processes in the plane*, volume 114 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2005.
- [16] G. F. Lawler, O. Schramm, and W. Werner. Conformal invariance of planar loop-erased random walks and uniform spanning trees. *Ann. Probab.*, 32(1B):939–995, 2004.
- [17] G. F. Lawler and W. Werner. The Brownian loop soup. *Probab. Theory Related Fields*, 128(4):565–588, 2004.
- [18] S. Rohde and O. Schramm. Basic properties of SLE. *Ann. of Math. (2)*, 161(2):883–924, 2005.
- [19] O. Schramm. Scaling limits of loop-erased random walks and uniform spanning trees. *Israel J. Math.*, 118:221–288, 2000.
- [20] O. Schramm. A percolation formula. *Electron. Comm. Probab.*, 6:115–120 (electronic), 2001.
- [21] S. Smirnov. Critical percolation in the plane. I. Conformal Invariance and Cardy’s formula II. Continuum scaling limit. *in preparation*, 2001.
- [22] W. Werner. Conformal restriction and related questions. *Lecture notes, ICMS Edinburgh, July 2003*, 2003.
- [23] W. Werner. Girsanov’s transformation for $SLE(\kappa, \rho)$ processes, intersection exponents and hiding exponents. *Ann. Fac. Sci. Toulouse Math. (6)*, 13(1):121–147, 2004.
- [24] W. Werner. Random planar curves and Schramm-Loewner evolutions. In *Lectures on probability theory and statistics*, volume 1840 of *Lecture Notes in Math.*, pages 107–195. Springer, Berlin, 2004.
- [25] D. B. Wilson. Generating random spanning trees more quickly than the cover time. In *Proceedings of the Twenty-eighth Annual ACM Symposium on the Theory of Computing (Philadelphia, PA, 1996)*, pages 296–303, New York, 1996. ACM.
- [26] D. Zhan. Stochastic Loewner Evolution in doubly connected domains. *Probab. Theory Related Fields*, 2003.
- [27] D. Zhan. Random Loewner chains in Riemann surfaces, 2004.

Courant Institute
 251 Mercer St., New York NY 10012
 dubedat@cims.nyu.edu